


Transcranial direct current stimulation beneficial in BED

May 17 2016



(HealthDay)—Transcranial direct current stimulation (tDCS) is

beneficial in binge-eating disorder (BED), with decreased cravings for sweets, savory proteins, and all foods, according to a study published online May 9 in the *International Journal of Eating Disorders*.

Emilee E. Burgess, from the University of Alabama at Birmingham, and colleagues examined the effect of tDCS on [food](#) craving, intake, binge-eating desire, and binge-eating frequency in 30 adults with BED or subthreshold BED. Participants received a 20-minute 2 mA session of tDCS targeting the [dorsolateral prefrontal cortex](#) and a sham session.

The researchers found that, compared with sham, tDCS correlated with decreased craving for sweets, savory proteins, and an all-foods category, with the strongest reductions seen in men (P food intake, respectively).

"This proof-of-concept study is the first to find ameliorating effects of tDCS in BED," the authors write. "The results support investigation of repeated tDCS as a safe and noninvasive treatment adjunct for BED."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Transcranial direct current stimulation beneficial in BED (2016, May 17) retrieved 30 April 2024 from
<https://medicalxpress.com/news/2016-05-transcranial-current-beneficial-bed.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
