


# Zika virus: Optimized tests for reliable diagnosis

May 12 2016

---



Transmission electron micrograph of Zika Virus particles (red). Credit: cdc/Cynthia Goldsmith

DZIF scientists from the University of Bonn have shown that not all conventional Zika virus molecular diagnostic tests for are sufficiently reliable. They developed optimised assays and a control for quantifying

viruses in blood and urine.

Diagnosing Zika viruses reliably is of major importance—for both patients and further research on the spread of the virus. The Zika virus is currently spreading in Central and South America; over one million people have become infected since spring 2015. Currently, acute infections are predominantly confirmed by determining the virus's genetic information in blood and urine. Six tests developed prior to the outbreak are currently being used in Central and South America. The problem is their reliability: can they also detect viral RNA in very low viral concentrations? How sensitive are they to different Zika virus strains and are their results comparable to each other? To date, these questions have not been fully clarified. Scientists therefore have concerns that numerous infections with the virus are going undetected.

## **The reliability problem**

"It is particularly important for pregnant women to reliably know whether they have Zika viruses in their blood or not," explains Prof Felix Drexler who, together with Prof Christian Drosten and his team from the University of Bonn, has now carefully examined the existing tests. Over the last months, it has been confirmed that a Zika virus infection during pregnancy can cause foetal brain malformations.

## **A comparison of all PCR tests**

In order to eliminate diagnostic uncertainties, the DZIF scientists initially tested the commonly used tests for sensitivity at the partner site Bonn-Cologne. All tests investigate viral RNA (ribonucleic acid), i.e. the hereditary information of the pathogen. The tests use the so-called polymerase chain reaction (PCR), a conventional method for detecting nucleic acid. The tests especially differ in that they examine different

regions of the viral gene. PCR tests are suitable for early virus detection in the first weeks after the onset of symptoms. Serological tests, which determine the antibodies produced by the infected person, are recommended for use after the eighth day.

## Results confirm concerns

The Zika assay comparison results confirm the researchers' concerns: some of the testing systems were not sensitive enough to detect low amounts of viruses. Additionally, not all [virus](#) strains are detected uniformly across the testing systems. Comparability between the assays is limited. The researchers assume that depending on the testing system, 20 to 80 percent of the patients may get an incorrect diagnosis, if serological testing methods are not used for further diagnosis.

## New testing systems for everyone

The researchers from Bonn consequently developed two new optimised PCR tests. Besides this, they have developed and made available a control which not only validates each [test](#), but also quantifies the viral RNA in the blood and urine. The so-called "calibrator" used for this purpose is synthetically constructed RNA which contains all the different viral RNA target zones used in the different conventional tests. The test protocols and the calibrator can be ordered worldwide free of charge.

"With our study, we especially wanted to call attention to the fact that a negative PCR test is not necessarily reliable," Drexler explains. The researchers have already made their results freely available prior to publication on the World Health Organisation (WHO) server. In an outbreak situation like this, all parties involved should exchange data as early as possible and have access to the best diagnostic tools.

The Bonn group led by Drexler and Drosten developed the globally used, standardised test for the MERS pathogen. At the DZIF, they are well-equipped for detecting newly emerging viruses. The Zika diagnostics project was supported by the DZIF and the European Union.

**More information:** Victor M. Corman et al, Clinical comparison, standardization and optimization of Zika virus molecular detection, *Bulletin of the World Health Organization* (2016). [DOI: 10.2471/BLT.16.175950](https://doi.org/10.2471/BLT.16.175950)

Provided by German Center for Infection Research

Citation: Zika virus: Optimized tests for reliable diagnosis (2016, May 12) retrieved 20 March 2024 from <https://medicalxpress.com/news/2016-05-zika-virus-optimized-reliable-diagnosis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--