

AAP: doctors should screen teens for suicide risk factors

June 29 2016



(HealthDay)—Suicide is the second leading cause of death among U.S.

teens, and health care providers should screen teen patients for suicide risks, according to a report published online June 27 in *Pediatrics*.

Suicide was the third leading cause of [teen](#) deaths in 2007, but has since passed homicide to become second. Unintentional injuries such as drug overdoses and vehicle crashes are the leading cause of teen deaths.

Teen suicide rates may have increased due to factors such as stress and anger triggered by electronic media, and a reluctance to use antidepressants, according to report author Benjamin Shain, M.D., Ph.D., head of the Division of Child and Adolescent Psychiatry at the NorthShore University HealthSystem in Chicago, *USA Today* reported.

Teen [suicide](#) risk factors that physicians should screen for include a history of physical or sexual abuse, bullying, substance abuse, mood disorders, and being lesbian, gay, or bisexual, according to the American Academy of Pediatrics.

More information: [Full Text](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: AAP: doctors should screen teens for suicide risk factors (2016, June 29) retrieved 17 April 2024 from <https://medicalxpress.com/news/2016-06-aap-doctors-screen-teens-suicide.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|