

# Eliminating blood test may increase availability of donor hearts

June 21 2016

---

A blood test that leads to the rejection of donor hearts may be unnecessary in predicting whether a heart transplant will succeed or fail, according to new research in *Circulation: Heart Failure*, an American Heart Association journal.

"Heart transplantation is an incredible therapy for patients with end-stage heart failure, but there are only 2,000-2,400 transplants each year. A lot of focus has been on finding ways to sign up more people as organ donors, but there is also a problem in that only an average of one in three donor hearts are placed," said Snehal R. Patel, M.D., senior author of the study and assistant professor of medicine at Montefiore Medical Center, Albert Einstein College of Medicine, in the Bronx, New York.

Many transplant centers routinely reject hearts if the donor's [blood test](#) reveals elevated levels of troponin I, a protein found in heart muscle that enters the bloodstream when there is a heart attack or other [heart muscle damage](#). Donors with previous heart disease are automatically excluded.

"If the heart looks okay but troponin I is elevated, many centers will reject the organ out of concern that this marker indicates dysfunction of the heart that will become evident after the stress of the transplant process. We looked at whether this is true," Patel said.

In the largest study of its kind and the first to use a national registry, researchers examined the outcomes of 10,943 adult (at least 18 years of age) [heart transplant recipients](#) in the national database of the United

Network of Organ Sharing. Each [donor heart](#) had normal pumping function.

Researchers found:

- no association between donor troponin I levels and the likelihood of death within the first year after transplant;
- no significant difference in survival at 30 days, 1 year, 3 years, or 5 years regardless of donor troponin I levels;
- no difference in the proportion of transplant recipients who developed primary graft failure, a catastrophic event marked by severe loss of pumping function in the donor heart that occurs within the first 30 days after transplant; and
- no difference in the proportion who developed cardiac allograft vasculopathy, a rapid form of [heart](#) disease that can occur in donor hearts and limits their long-term success.

"Our research shows that [transplant centers](#) should not exclude donor hearts based solely on elevated troponin I if the organ is otherwise suitable. At our institution it has already changed how we evaluate donors, and I think this data will lead to changes nationwide," Patel said.

Provided by American Heart Association

Citation: Eliminating blood test may increase availability of donor hearts (2016, June 21)  
retrieved 3 May 2024 from

<https://medicalxpress.com/news/2016-06-blood-availability-donor-hearts.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--