


Contaminated gloves increase risks of cross-transmission of pathogens

June 20 2016



Credit: American Society for Microbiology

Research being presented at the ASM Microbe research meeting provides clear evidence that the gloves of healthcare workers contaminate hospital surfaces with bacteria. The researchers' data also suggest that types of bacteria may affect cross-transmission rates among

contaminated gloves and the hospital surfaces.

"Infection control is a priority for all hospitals to reduce the spread of HCAs," said study author Sae Otani, a master course student at Bunkyo Gakuin University, "Gloving is recommended as a barrier protection for [healthcare workers](#) to reduce the risk of contamination during contact with infectious sputum, urine and body fluids," she said. However, failure to change or remove contaminated gloves carries a high-risk of healthcare-associated pathogens transmission.

The researchers, led by Sae Otani, and Kazue Fujita, Nippon Medical School used common healthcare-associated pathogens in this study. Nitrile examination gloves were inoculated with the specific microorganism. After inoculation of [bacteria](#), contaminated gloves were touched to a sterilized polypropylene surface. Then, the number of viable bacteria on the polypropylene surface was quantified

Acinetobacter baumannii remained on the polypropylene surface. In contrast, *Escherichia coli*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* were not detected on the polypropylene surface. Antibiotic sensitivity of bacteria had no consistent effect on survival on the polypropylene surface.

"This study shows that contaminated gloves increase risks of cross-transmission of healthcare-associated pathogens among healthcare workers and in the environment," said Ms. Otani. In addition, the data also suggest that types of bacteria may affect cross-transmission rates among contaminated gloves and the hospital surfaces. *A. baumannii* are higher risk of transmission from the gloves to hospital surfaces than *E. coli*, *K. pneumoniae* and *P. aeruginosa* in this study. HCAs caused by gram-negative organisms such as *Acinetobacter* spp., *P. aeruginosa* and Enterobacteriaceae are a major concern.

"Proper glove use may decrease the risk of healthcare-associated infections and gloves should be carefully used and removed after use depending on the types of bacteria," said Ms. Otani.

Provided by American Society for Microbiology

Citation: Contaminated gloves increase risks of cross-transmission of pathogens (2016, June 20) retrieved 24 April 2024 from

<https://medicalxpress.com/news/2016-06-contaminated-gloves-cross-transmission-pathogens.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.