

Foot wrap offers alternative to medication for patients with restless legs syndrome

June 27 2016

Authors from Lake Erie Research Institute in Pennsylvania report an adjustable foot wrap caused to treat restless legs syndrome (RLS) is 1.4 times more effective than the standard pharmaceutical treatment. The pilot study published today in *The Journal of the American Osteopathic Association*.

The eight-week clinical trial involved 30 otherwise healthy adults with moderate to severe restless leg syndrome. Researchers studied Clinical Global Impression responses as well as the mean change in the International Restless Leg Syndrome Study Group Study Scale (IRLSSGS). A meta-analysis was then used to compare the RLS device with three historic studies of the medication ropinirole and a placebo.

Clinical Global Impression responses indicated significantly greater improvement with the RLS device (90 percent) compared with ropinirole (63 percent), the current standard dopamine therapy for RLS. Additionally, change in IRLSSGS score was significantly greater for the RLS device (17.22) compared to historic reports for ropinirole versus the placebo (12 versus 8.9 respectively). Patients using the RLS device also reported an 82 percent decrease in sleep loss.

The RLS device was designed to put adjustable targeted pressure on two muscles in the foot known to relax symptoms of RLS, the abductor hallucis and the flexor hallucis brevis. Researchers indicate that the pressure produced by the device may also stimulate a dopamine release, similar to massage therapy or acupressure.

"By putting pressure on specific muscles in the feet, we are able to create a response in the brain that relaxes the muscles activated during RLS," said Phyllis Kuhn, MS, PhD, and the study's lead researcher. "It's a near perfect example of the body regulating itself without drugs, many of which have the potential for significant adverse side effects."

Restless leg syndrome (RLS) is a neurologic disorder causing unpleasant sensations and an urge to move the legs when at rest. The sleep loss associated with RLS can cause extreme fatigue, anxiety and depression. According to the National Institute of Health, RLS may affect as many as 10 percent of the U.S. population, with more than nine million experiencing moderate to severe symptoms.

Until recently, potent drugs including opioids, depressants and dopamine agonists have been used to ease symptoms, but each of these is accompanied by negative side effects such as dizziness, nausea, vomiting and the added risk of addiction.

"Restless legs syndrome really erodes quality of life because it causes extreme fatigue for many patients. As an osteopathic physician, it's a challenge to balance the need to restore sleep while preventing additional harm from medication. These results show promise in otherwise healthy individuals for a nonpharmaceutical option that appears to have rather minor, temporary adverse effects for some users," said Rob Danoff, DO, an osteopathic family physician and program at Aria Health Care in Philadelphia.

Adverse effects were reported by seven patients in the study. The effects included pain (1), pins and needles sensation (2), irritability (3), spasm (1) and warm feet (1).

More information: *The Journal of the American Osteopathic Association*, jaoa.org/article.aspx?articleid=2531565 , DOI:

[10.7556/jaoa.2016.088](https://doi.org/10.7556/jaoa.2016.088)

Provided by American Osteopathic Association

Citation: Foot wrap offers alternative to medication for patients with restless legs syndrome (2016, June 27) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-06-foot-alternative-medication-patients-restless.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.