

Gastric bypass cuts hormonal responses to hypoglycemia

June 29 2016



(HealthDay)—Gastric bypass (GBP) surgery is associated with reduced



symptoms and neurohormonal responses to hypoglycemia, according to a study published online June 16 in *Diabetes*.

Noting that GBP patients have lowered glucose levels and frequent asymptomatic hypoglycemic episodes, Niclas Abrahamsson, M.D., Ph.D., from Uppsala University in Sweden, and colleagues examined symptoms and hormonal and autonomic nerve responses in patients exposed to hypoglycemia before and after GBP. Twelve obese patients without diabetes underwent hyperinsulinemic hypoglycemic clamp before and after GBP surgery.

The researchers found that after surgery the Edinburgh Hypoglycemia symptom delta scores during clamp were attenuated from 10.7 to 5.2. Marked reductions in glucagon, cortisol, catecholamine, and sympathetic nerve responses to hypoglycemia were seen after surgery. A delayed response was seen in growth hormones, but to a higher peak level. During hypoglycemia, glucagon-like peptide-1 and gastric inhibitory peptide increased, but to a lesser extent after surgery.

"GBP surgery causes a resetting of glucose homeostasis, which reduces symptoms and neurohormonal responses to hypoglycemia," the authors write. "Further studies should address the underlying mechanisms as well as their impact on the overall metabolic effects of GBP surgery."

The study was partially funded by Exodiab.

More information: <u>Full Text (subscription or payment may be required)</u>

Copyright © 2016 HealthDay. All rights reserved.

Citation: Gastric bypass cuts hormonal responses to hypoglycemia (2016, June 29) retrieved 19



April 2024 from

https://medicalxpress.com/news/2016-06-gastric-bypass-hormonal-responses-hypoglycemia.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.