


# HIV infection negatively affects bone acquisition

June 15 2016

---



(HealthDay)—HIV infection with T cell activation is associated with

lower bone mineral density (BMD) and stiffness, according to a study published online June 10 in the *Journal of Bone and Mineral Research*.

John S. Manavalan, M.D., from the Columbia University Medical Center in New York City, and colleagues used dual X-ray absorptiometry (DXA) and high-resolution peripheral quantitative computed tomography (HRpQCT) to characterize BMD and microarchitecture in 38 HIV-infected men on antiretroviral therapy (18 perinatally-infected and 20 adolescence-infected) and 20 uninfected men aged 20 to 25 years.

The researchers found that, compared with uninfected men, HIV-infected men had lower DXA derived areal BMD Z-scores and HRpQCT derived volumetric BMD measures. HIV-infected men had higher proportions of activated and senescent CD4+ and CD8+ T cells compared with uninfected men. HIV-infected men had a lower percentage of circulating osteogenic precursor (COP) cells than uninfected men ( $0.19 \pm 0.02$  versus  $0.43 \pm 0.06$  percent; P stiffness and a lower proportion of COP [cells](#).

"T cell activation with HIV-infection was associated with decreased numbers of osteogenic precursors as well as lower peak bone mass and [bone](#) strength," the authors write.

One author disclosed financial ties to Gilead and AbbVie.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: HIV infection negatively affects bone acquisition (2016, June 15) retrieved 19 April

2024 from <https://medicalxpress.com/news/2016-06-hiv-infection-negatively-affects-bone.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.