

# International trial changing standard of care for advanced breast cancer

June 2 2016

---

Surgery to remove the primary tumor in women diagnosed with stage IV breast cancer, followed by the standard combination of therapies, adds months to the patients' lives, compared with standard therapy alone, an international clinical trial led by a University of Pittsburgh Cancer Institute (UPCI) professor revealed.

The results of the phase III randomized, controlled trial will be presented Saturday at the American Society of Clinical Oncology (ASCO) annual meeting in Chicago. The study was selected for the society's "Best of ASCO," an effort to condense the research "most relevant and significant to oncology" into a two-day program to increase global access to cutting-edge science.

"Our findings will change the standard of care for [women](#) newly diagnosed with stage IV [breast cancer](#)," said principal investigator Atilla Soran, M.D., M.P.H., clinical professor of surgery, University of Pittsburgh School of Medicine, and breast surgical oncologist with UPMC CancerCenter. "We've shown that surgery to remove the primary tumor—either through lumpectomy or mastectomy—followed by standard therapy, is beneficial over no surgery."

Dr. Soran began the trial in 2007, ultimately recruiting a total of 274 women newly diagnosed with stage IV breast cancer from 25 institutions. Half the women received standard therapy, which avoids surgery and consists of a combination of chemotherapy, hormonal therapy and targeted therapy, while the other half first had surgery to

remove their primary breast tumor, followed by the standard therapy.

At about 40 months after diagnosis, the women who received the surgery plus standard therapy lived an average of nine months longer than their counterparts who received standard therapy alone. Nearly 42 percent of the women who received surgery lived to five years after diagnosis, compared with less than 25 percent of the women who did not receive surgery.

The trial also showed that [surgery](#) in younger women with less aggressive cancers resulted in longer average survival than in women with more aggressive cancers that had spread to the liver or lungs.

"Our thinking is similar to how you might approach a battle against two enemies," said Dr. Soran. "First you quickly dispatch one army—the primary tumor—leaving you to concentrate all your efforts on battling the second army—any remaining cancer."

Provided by University of Pittsburgh Schools of the Health Sciences

Citation: International trial changing standard of care for advanced breast cancer (2016, June 2) retrieved 10 April 2024 from

<https://medicalxpress.com/news/2016-06-international-trial-standard-advanced-breast.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--