


Materials in vitreous seen in about half of phaco cases

June 23 2016



(HealthDay)—Materials in the vitreous (MIV) are identified in

approximately half of phacoemulsification cataract surgery (phaco) cases, in Berger's space or the anterior vitreous (AV), according to a study published online June 17 in *Clinical & Experimental Ophthalmology*.

Andrew W. Kam, from the Prince of Wales Hospital in Sydney, and colleagues conducted a prospective observational study involving 767 consecutive phaco [cases](#) to identify and classify MIV.

The researchers observed MIV in 50.3 percent of cases (386 eyes), either in Berger's space or the AV. Most MIV was suspected to be lens material (46.5 percent of all cases). In 9.8 and 1.7 percent of cases, respectively, pigment and ophthalmic viscoelastic devices were seen in the AV. Positive predictors for the presence of MIV included higher nuclear sclerosis grade, male gender, and older age. The MIV was identified as lens material in histological assessment with light microscopy and birefringence techniques.

"MIV were seen in 50.3 percent of phaco cases," the authors write. "It has been histologically demonstrated that [lens](#) materials can be introduced into the AV during phaco."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Materials in vitreous seen in about half of phaco cases (2016, June 23) retrieved 17 April 2024 from <https://medicalxpress.com/news/2016-06-materials-vitreous-phaco-cases.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.