

Opinions vary on management of subclinical hypothyroidism

June 7 2016



(HealthDay)—Physicians' opinions on management of subclinical



hypothyroidism vary, according to a Beyond the Guidelines article published online June 7 in the *Annals of Internal Medicine*.

In an effort to examine approaches to management of subclinical hypothyroidism, Risa B. Burns, M.D., M.P.H., from the Beth Israel Deaconess Medical Center in Boston, and colleagues discuss the case of a 60-year-old woman (Ms. C) with a positive family history of thyroid disease, elevated thyroid stimulating hormone (TSH), and symptoms compatible with hypothyroidism.

The author in favor of treatment would recommend levothyroxine treatment based on elevated TSH and available evidence, including hyperlipidemia. With low-dose treatment and careful monitoring, treatment would be unlikely to result in harm. Treatment could be stopped if the patient did not feel better and cholesterol was not improved. The author against treatment notes that the patient's degree of thyroid dysfunction would not increase the risk of heart disease, and that the goals of weight loss and symptom relief would be unlikely to be achieved with medication. If the patient did start treatment, the dose should be 25 to 50 mcg to avoid overtreatment, and careful monitoring should be conducted. In the event her symptoms did not improve, treatment should be stopped.

"While we await further data, both our discussants concur with shared decision-making so that Ms. C makes a fully informed decision," the authors write.

More information: <u>Full Text (subscription or payment may be required)</u>

Copyright © 2016 HealthDay. All rights reserved.



Citation: Opinions vary on management of subclinical hypothyroidism (2016, June 7) retrieved 19 April 2024 from

https://medicalxpress.com/news/2016-06-opinions-vary-subclinical-hypothyroidism.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.