


# Physical activity in dementia

June 8 2016, by Ríona Mc Ardle



This figure shows the amount of time each group spent physically active over a 24 hour period. The healthy control group engages in the largest amount of activity, with the institutionalized dementia group engaging in significantly less. Credit: Van Alphen, et. al. <http://dx.doi.org/10.1371/journal.pone.0152457>

It has been recognised for over 60 years that regular physical activity provides health benefits. Our bodies are designed to be stimulated by physical exertion. Fitness is associated with lower all-cause mortality, along with improved sleep, quality of life, social life and reduced stress. On the other hand, physical inactivity predicts obesity, diabetes, cardiovascular disease, mental health issues, reduced quality of life and


overall mortality. The evidence is clear: we need to get moving.

In addition to health outcomes, cognition and physical activity are intrinsically linked. This is increasingly true within the senior population. More physically active older adults show less cognitive decline than those who are not active. Older adults participating in physical activity programs often demonstrate improvements in their cognitive abilities – particularly with attention and executive function. In frailer older adults with many health problems, physical activity interventions facilitate better cognition, physical capacity and quality of life. Regular physical activity has also been associated with reduced risk of cognitive decline and development of dementia. Many hypotheses have been postulated regarding the role of physical activity in staving off [cognitive impairment](#). It may protect cardiovascular changes in the brain by maintaining cerebral blood flow, facilitate the growth of new neurons or relieve stress and therefore improve cognitive function.

But what does physical activity look like in the cognitively impaired population – specifically within dementia? A recent study in *PLOS ONE* aimed to quantify and characterize physical activity in dementia [patients](#) objectively. While previous studies report that dementia patients are less physically active than [older adults](#) without cognitive impairment, the majority of these findings come from self-reported measures. These are often not as reliable or accurate as they are subjective, and may be impacted by memory recall problems. Van Alphen and colleagues chose to use the [Actiwatch](#), a wrist-worn accelerometer, to collect objective data on the frequency, duration and intensity of daily physical activity in dementia patients. One hundred and twenty dementia patients were included in the study, recruited from 13 nursing homes and four day-care centers, along with 26 older adult controls. All participants were required to wear the Actiwatch for at least six days, 24 hours a day.

The researchers were interested in the differences in [physical activity](#)

levels between community-dwelling dementia patients and institutionalized patients. To this end, daily physical activity levels were significantly different between institutionalized patients and community-dwelling patients and controls, with institutionalized patients engaging in physical activity 23.5% less than community-dwelling patients and 40% less than controls. This is important to note as infrastructures and activities could be implemented in care homes to encourage physical activity.



This figure demonstrates the amount of time the different groups spend

sedentary compared to other levels of physical intensity. The first bars (0-99) represent sedentary time, while the rest of the bars show a gradual climb in activity intensity. Credit: Van Alphen, et. al.

<http://dx.doi.org/10.1371/journal.pone.0152457>

Additionally, institutionalized patients spent 72% of their day in a sedentary state; 9.3% more than community-dwelling patients and 19% more than controls. These results remain significant after controlling for age, cognitive abilities and use of walking aids. While there were no significant differences in amount of physical activity between Alzheimer's disease patients and non-Alzheimer's disease patients (encompassing dementia with Lewy bodies, Parkinson's disease with dementia, vascular dementia and frontotemporal dementia), Alzheimer's disease patients did spend significantly less time (6.9%) in a sedentary state. Van Alphen and colleagues concluded that both institutionalized and community-dwelling dementia patients remained sedentary for the majority of the day and performed lower intensity activities than that of controls.

But why is it important to know about physical activity levels in dementia? The dangers of physical inactivity were highlighted earlier; for [dementia patients](#), the ability to carry out everyday activities may be compromised, along with reduced quality of life and further cognitive decline. Studies have demonstrated that encouraging active participation in functional tasks – such as dressing and washing – and attending dance or exercise classes – can improve physical capacity and reduce fall risk in moderate-to-severe dementia. Bi-weekly exercise programs are associated with slower rates of cognitive and functional decline in dementia. Individuals with mild cognitive impairment – a pre-dementia stage – undergoing an aerobic fitness intervention demonstrate significant improvements in cognition, particularly executive function.

Encouraging physical activity in this population holds a range of benefits – from improved mood, function, cognition and fitness to reduced stress and improved quality of life for their caregiver.

There is still much work to be done. Studies need to recruit larger samples of patients with non-Alzheimer's disease dementias to assess the differences across dementia subtypes. Type of activity should be examined; most accelerometers can provide data about frequency and duration of movement but fail to automatically specify what activities are being undertaken. Individual differences in activity type may predict different health outcomes. As Van Alphen and colleagues reported, [dementia](#) groups are more likely to engage in low-intensity activities such as walking or household chores. These activities are beneficial for cardiorespiratory capacity and body composition, and exercise interventions could be tailored to suit this preference. The effects of vigorous – such as sprinting or spinning – and low-intensity activities should also be compared. This is an area ripe for investigation and may be beneficial for the development of physical interventions.

Increasing physical activity levels is a global challenge and one that is continuously being addressed. [The World Health Organization \(WHO\)](#) recommends 30 minutes of moderate-to- vigorous exercise a day, which can be achieved by walking, gym classes and activities, sports or physical hobbies such as gardening. Governments develop policies, guidelines and initiatives to encourage people to participate in physical exercises, such as [Let's Move](#). National campaigns – such as [This Girl Can](#) – promote a sense of unity in keeping active. Free exercise programs – such as Shape Up New York – allow people to engage in a range of fitness classes, from aerobics to Zumba. These all endorse the message to get up, get going and get active! Still 20% of individuals older than 15 years do not meet these physical activity guidelines. Therefore, I believe our approach to the promotion and facilitation of [physical activity](#) needs to change. Most people associate the prescription of getting active with

problems like obesity and heart disease. More education on how active individuals incur health benefits beyond physical health is also needed.

Exercise is often portrayed as inaccessible without expensive gym facilities and a grueling experience for those who do take part – think of all those films depicting the embarrassment of being picked last in gym class. Facilities allowing engagement in moderate-intensity exercise – such as walking, cycling and running – should be built; fun and simple ways to get involved should be endorsed. Physical activity is an inexpensive, easily-accessible approach to improve health and overall well-being – and we should be taking full advantage of that.

**More information:** Helena J. M. van Alphen et al. Older Adults with Dementia Are Sedentary for Most of the Day, *PLOS ONE* (2016). [DOI: 10.1371/journal.pone.0152457](https://doi.org/10.1371/journal.pone.0152457)

Helena J. M. van Alphen et al. Older Adults with Dementia Are Sedentary for Most of the Day, *PLOS ONE* (2016). [DOI: 10.1371/journal.pone.0152457](https://doi.org/10.1371/journal.pone.0152457)

Renée F. A. G. de Bruijn et al. The association between physical activity and dementia in an elderly population: the Rotterdam Study, *European Journal of Epidemiology* (2013). [DOI: 10.1007/s10654-013-9773-3](https://doi.org/10.1007/s10654-013-9773-3)

Samuel C. Dumith et al. Worldwide prevalence of physical inactivity and its association with human development index in 76 countries, *Preventive Medicine* (2011). [DOI: 10.1016/j.ypmed.2011.02.017](https://doi.org/10.1016/j.ypmed.2011.02.017)

Laura D. Baker et al. Effects of Aerobic Exercise on Mild Cognitive Impairment, *Archives of Neurology* (2010). [DOI: 10.1001/archneurol.2009.307](https://doi.org/10.1001/archneurol.2009.307)

Nicola T. Lautenschlager et al. Physical Activity and Mild Cognitive

Impairment and Alzheimer's Disease, *Current Neurology and Neuroscience Reports* (2010). [DOI: 10.1007/s11910-010-0121-7](https://doi.org/10.1007/s11910-010-0121-7)

E. Galik et al. Optimizing Function and Physical Activity Among Nursing Home Residents With Dementia: Testing the Impact of Function-Focused Care, *The Gerontologist* (2013). [DOI: 10.1093/geront/gnt108](https://doi.org/10.1093/geront/gnt108)

Physical activity for improving cognition in older people with mild cognitive impairment [DOI: 10.1002/14651858.CD008198](https://doi.org/10.1002/14651858.CD008198)

The Role Of Physical Activity On The Prevention Of Cognitive Impairment. [www.ncbi.nlm.nih.gov/pmc/articles/PMC4811348/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4811348/)

Louis Bherer et al. A Review of the Effects of Physical Activity and Exercise on Cognitive and Brain Functions in Older Adults, [www.w3.org/1999/xhtml](http://www.w3.org/1999/xhtml)">*Journal of Aging Research* (2013). [DOI: 10.1155/2013/657508](https://doi.org/10.1155/2013/657508)

Pedro C Hallal et al. Physical activity: more of the same is not enough, *The Lancet* (2012). [DOI: 10.1016/S0140-6736\(12\)61027-7](https://doi.org/10.1016/S0140-6736(12)61027-7)

Pamela Das et al. Rethinking our approach to physical activity, *The Lancet* (2012). [DOI: 10.1016/S0140-6736\(12\)61024-1](https://doi.org/10.1016/S0140-6736(12)61024-1)

*This story is republished courtesy of PLOS Blogs: [blogs.plos.org](https://blogs.plos.org).*

Provided by Public Library of Science

Citation: Physical activity in dementia (2016, June 8) retrieved 24 April 2024 from <https://medicalxpress.com/news/2016-06-physical-dementia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.