

Increased prevalence of HPV for men with phimosis

June 16 2016



(HealthDay)—Men with phimosis have increased prevalence of human

papillomavirus (HPV) and high-risk HPV genotypes compared with asymptomatic men, according to a study published in the June issue of *The Journal of Urology*.

Larissa A. Afonso, Ph.D., from the Universidade Federal Fluminense in Rio de Janeiro, and colleagues examined the incidence of HPV [infection](#) in 110 asymptomatic men and 30 [patients](#) who underwent circumcision due to phimosis. DNA was extracted from swabbed samples from asymptomatic men and from foreskin samples obtained at circumcision in men with phimosis.

The researchers found that 46.66 percent of patients with phimosis had HPV, of whom 50 percent had high-risk HPV genotypes. Overall, 16.36 percent of the asymptomatic cases were HPV positive; high-risk HPV was seen in only one sample. Patients presenting with phimosis had a significantly higher rate of HPV genital infection than asymptomatic men ($P = 0.00167$). Patients with phimosis had significantly higher prevalence of high-risk HPV genotypes ($P = 0.004$).

"We found a robust association between phimosis and the genital HPV prevalence in men and a significant frequency of high-risk HPV," the authors write. "Other studies are needed to investigate the occurrence of factors that can increase the incidence of penile carcinoma and determine its impact on female genital infection in cervical cancer."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Increased prevalence of HPV for men with phimosis (2016, June 16) retrieved 10 April 2024 from <https://medicalxpress.com/news/2016-06-prevalence-hpv-men-phimosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.