


PTSD may negatively affect sex life satisfaction in male and female veterans

June 3 2016



Regions of the brain associated with stress and posttraumatic stress disorder. Credit: National Institutes of Health

New research reveals that posttraumatic stress disorder (PTSD) was a strong, negative predictor of sexual satisfaction in both male and female veterans who returned from warzones in recent Iraq and Afghanistan conflicts.



For the *Journal of Traumatic Stress* study, investigators assessed 1581 males and females from the Veterans After-Discharge Longitudinal Registry (VALOR), a gender-balanced registry of VA healthcare-seeking <u>veterans</u> with and without PTSD.

Sexual activity rates were similarly high in male veterans with and without a PTSD diagnosis, whereas PTSD status was negatively associated with sexual activity rates in <u>female veterans</u>.

"Future analysis in the longitudinal cohort of the VALOR registry is uniquely suited to answer questions regarding the influence of PTSD treatment on sexual dysfunction, activity and satisfaction," the authors wrote.

More information: Benjamin N. Breyer et al, Sexual Health in Male and Female Iraq and Afghanistan U. S. War Veterans With and Without PTSD: Findings From the VALOR Cohort, *Journal of Traumatic Stress* (2016). DOI: 10.1002/jts.22097

Provided by Wiley

Citation: PTSD may negatively affect sex life satisfaction in male and female veterans (2016, June 3) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2016-06-ptsd-negatively-affect-sex-life.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.