

Study reveals high rate of disordered eating in young Australian women

June 28 2016



Credit: University of Queensland

Up to one-third of young Australian women experience episodes of binge or overeating, with socially disadvantaged women at greater than average risk.

Researchers from The University of Queensland, Stockholm University and the Karolinska Institutet in Sweden have found that four per cent of women aged 18 to 23 reported symptoms of bulimia nervosa, but the incidence of milder [eating disorders](#) was much greater.

UQ School of Public Health's Professor Gita Mishra said 17 per cent of women reported episodes of [overeating](#), 16 per cent reported [binge eating](#) and 10 per cent reported compensatory behaviours such as vomiting, use of laxatives or diuretics and fasting.

"The results highlight the large burden of both transient and persistent milder and undiagnosed forms of disordered and overeating in this age group," Professor Mishra said.

The study analysed data from more than 6800 participants in the Australian Longitudinal Study on Women's Health (ALSWH), building on earlier work on social and early life causes of eating disorders at the Centre for Health Equity Studies (CHESS) in Sweden.

Professor Ilona Koupil from CHESS said the research showed there was an increased risk of binge eating or overeating among socially disadvantaged women.

"There was also increased risk among young women who reported smoking and binge drinking, suggesting a possible overlap between substance abuse and eating disorders," Professor Koupil said.

The researchers believe social patterns could be used to identify at-risk groups and those in need of early diagnosis and secondary prevention.

"We were intrigued to see a higher risk of binge eating and bulimia nervosa among women of European origin and in those who had been overweight or obese in childhood," Professor Koupil said.

"We hope our results prompt further efforts to monitor prevalence of these common disorders among young [women](#) in Australia and overseas."

The researchers said more investigation was needed into long-term [health](#) outcomes and consequences of disordered eating.

The study is published in *Public Health Nutrition*.

More information: Ilona Koupil et al. Social patterning of overeating, binge eating, compensatory behaviours and symptoms of bulimia nervosa in young adult women: results from the Australian Longitudinal Study on Women's Health, *Public Health Nutrition* (2016). [DOI: 10.1017/S1368980016001440](#)

Provided by University of Queensland

Citation: Study reveals high rate of disordered eating in young Australian women (2016, June 28) retrieved 1 May 2024 from <https://medicalxpress.com/news/2016-06-reveals-high-disordered-young-australian.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--