

Scientists detect gray matter changes in brains of teenagers with type 2 diabetes

June 14 2016



Credit: public domain

Teenagers with type 2 diabetes have significant changes in total brain gray matter volume and in regions of gray matter involved in seeing and hearing, memory, emotions, speech, decision making, and self-control.

Researchers at Cincinnati Children's Hospital Medical Center discovered that these teens have six regions with significantly less gray matter, and three with significantly more. They found a relationship between less [gray matter](#) volume in the brain and the ability to pronounce and sound

out unfamiliar words.

"Previous studies suggested that youth with type 2 [diabetes](#) have changes in brain structure and poorer cognitive function scores compared to their peers," says Amy Sanghavi Shah, MD, a physician-scientist in the division of Endocrinology at Cincinnati Children's and senior author of the study. "Total and regional brain volume had not been assessed comprehensively until now. We also sought to determine if the findings we found here could explain poorer cognitive scores."

The study will be presented at 10:45 am Central time Tuesday, June 14, 2016 at the American Diabetes Association's Scientific Sessions in New Orleans. Jacob Redel, MD, a fellow in the division of Endocrinology at Cincinnati Children's and lead author of the study, will present the findings at the ADA Presidents Oral Session.

The researchers studied 20 teens with type 2 diabetes and compared them to 20 teens without type 2 who were similar in age, race and sex. All participants in the study had high-resolution MRIs. Neither group had prior neurological or psychological disease or prior abnormal MRIs.

"Our results do not show cause and effect," says Dr. Redel. "We don't know if the changes we found are the direct result of diabetes, but studies in adults with type 2 diabetes with longer duration of disease also show brain volume differences, brain vascular changes and cognitive decline. However, our findings suggest that preventing type 2 diabetes in adolescents is important to prevent possible complications in the future."

The researchers are considering another study adding participants who are obese without type 2 diabetes and a larger number of participants. This would allow them to examine whether differences found on MRI are more related to obesity or [high blood sugar](#) seen in diabetes and test whether other cognitive domains, such as memory, language, intellect

and attention, are affected by [brain volume](#) differences.

Provided by Cincinnati Children's Hospital Medical Center

Citation: Scientists detect gray matter changes in brains of teenagers with type 2 diabetes (2016, June 14) retrieved 27 April 2024 from <https://medicalxpress.com/news/2016-06-scientists-gray-brains-teenagers-diabetes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--