


Southern Europe risks Zika outbreaks this summer

June 10 2016



Risk areas for August. Reproductive number (R_0) explains the estimated epidemic growth rate of Zika. At levels below 1 epidemics die out. At higher values the epidemic grows exponentially. Credit: *EBioMedicine*

Established *Aedes-mosquito* population could spread the Zika virus in Europe this summer if infected travelers introduce the virus. An analysis of temperatures, vectorial capacity, basic reproductive number (R0), and air traveler flows suggests parts of Southern Europe may be at risk for Zika outbreaks between June and August. This according to a study, led by Umeå University researchers in Sweden and published in the journal *EBioMedicine*.


"We know warm climates create the kind of conditions suitable for mosquito-borne illnesses to spread," says Joacim Rocklöv, researcher at Umeå University's Unit for Epidemiology and Global Health and co-author of the article.

"Vectorial capacity depends on a number of parameters but in general, warmer temperatures increase the rate in which the female mosquitos bite, the mosquito virus reproduction, and their virus transmission [risk](#). The presence of established *Aedes mosquito* populations, the warmer climate and the coinciding peak flow of air travelers into Europe, is a triage making Southern Europe fertile ground for Zika."



Following a similar epidemiological study conducted on the similar dengue virus, the group of researchers led by Joacim Rocklöv at Umeå University, used a temperature dependent computer model to predict Zika virus infection risks for Europe. The research exploration was undertaken in close collaboration with the European Centre for Disease Prevention and Control (ECDC). In the analysis, the researchers overlaid data on monthly flows of airline travelers arriving in European cities from Zika-affected areas, data on month-by-month estimates of virus infection reproduction capabilities of *Aedes-mosquito* populations in Europe, and human population data within the areas where mosquito-borne transmission of the Zika virus could be possible.

JUNE

MONTHLY VOLUME OF TRAVELLERS




REPRODUCTION NUMBER (R_J)






JULY

MONTHLY VOLUME OF TRAVELLERS




REPRODUCTION NUMBER (R_J)






AUGUST

MONTHLY VOLUME OF TRAVELLERS



REPRODUCTION NUMBER (R_J)



Risk areas for Zika in June-August. Reproductive number (R_0) explains the estimated epidemic growth rate of Zika. At levels below 1 epidemics die out. At higher values the epidemic grows exponentially. Credit: *EBioMedicine*

The main findings, presented in *EBioMedicine*, a journal initiative integrating the *Lancet* and *Cell*, are:

- The risk of mosquito-borne transmission of Zika virus is estimated to peak between June and August in parts of Southern Europe (see map)
- The peak flow of air travelers from regions of the Americas affected by the Zika virus coincides with the peak in the *Aedes-mosquitos* capacity to transmit the virus.
- The findings could help European public health officials to identify locations and times where the risk for Zika is heightened.

The risk assessment assumes that European *Aedes-mosquitos* have the same potential to spread the Zika virus as their South-, Middle- and North American counterparts.

Earlier research has shown that increasing temperatures will enlarge Europe's seasonal window for the potential spread of mosquito-borne viral disease and expand the geographic areas at risk for epidemics to include large parts of Europe. The threat includes tropical and sub-tropical viruses such as Zika and Dengue.

The *Aedes mosquitos*—*Aedes aegypti* and *Aedes albopictus*—are largely responsible for the transmission of the Zika [virus](#). Both *Aedes* mosquitos

are likely to become a fixture in Europe. Historically, *Aedes* mosquitos were present in many European countries during the first half of the 1900s. *Aedes aegypti* has recently been documented in Russia and Georgia. And current surveillence indicate that *Aedes albopictus* are present in much of Southern Europe and as far north as the Netherlands.

More information: Assessing Seasonal Risks for the Introduction and Mosquito-borne Spread of Zika Virus in Europe, [DOI: 10.1016/j.ebiom.2016.06.009](https://doi.org/10.1016/j.ebiom.2016.06.009) , www.ebiomedicine.com/article/S2352-3964%2816%2930253-5/abstract

Provided by Umea University

Citation: Southern Europe risks Zika outbreaks this summer (2016, June 10) retrieved 11 May 2024 from <https://medicalxpress.com/news/2016-06-southern-europe-zika-outbreaks-summer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--