

Special properties of pneumococci affect their ability to cause meningitis

June 28 2016



Credit: AI-generated image (disclaimer)

Structures on the surface of pneumococci determine the ability of these bacteria to enter the brain and cause severe infections, according to a paper published in the *Journal of Clinical Investigation* by researchers at Karolinska Institutet. Their findings suggest that certain pneumococci that are small in size, and which have a special protein on their surface,



can more easily pass through the blood-brain barrier and cause meningitis that may be lethal.

Pneumococci are known for causing infections in the respiratory tract such as sinusitis, otitis and pneumonia. They are also a common cause of severe diseases such as <u>blood</u> poisoning (septicemia) and meningitis. But how pneumococci enter the <u>brain</u> through the blood-brain barrier – the protective network of capillaries that prevents drugs and potentially harmful cells (e.g. immune cells, bacteria and viruses) from passing from the blood to the brain – is unclear.

Structures on surface

The current study, which was led by professor Birgitta Henriques-Normark, demonstrates that structures on the surface of the bacteria, socalled pili, affect how they can pass through the blood -brain barrier and infect the brain tissue. The researchers examined more closely one of the pilus proteins, RrgA, which they have shown is important for how pneumococci can bind to cells in the lung in previous studies. They now discovered that RrgA facilitates the passage of pneumococci through the blood-brain barrier.

Pneumococci have a round ovoid form, and when they proliferate in the blood stream they usually appear in pairs or in short chains. However, the researchers have now found that a small proportion of pneumococci with pili in the blood, less than five per cent, can be found as individual spherical cocci. And it is these single cocci that promote penetration of the bacteria through the blood-brain barrier. Moreover, the researchers found that pneumococci with pili that infected brains were spherical, single cocci that did not express the protein DivIVA, which is involved in cell division and responsible for the oval-shaped form of pneumococci.



"Our data suggests that <u>pneumococci</u> that have the pilus protein RrgA, and that present as single cocci and are small in size, more easily pass from the blood into the brain, where they can cause potentially lifethreatening meningitis," says principal investigator Professor Henriques-Normark at the Department of Microbiology, Tumor and Cell Biology.

Serious consequences

Meningitis is an acute, suppurative infection of the brain membranes and the spaces in between them. Meningitis is most common in children and adults over the age of 60, and can trigger symptoms such as fever, headaches, stiff neck, insomnia, vomiting, spasms and coma. The disease can have serious long-term consequences and is often fatal.

More information: Federico Iovino et al. Pneumococcal meningitis is promoted by single cocci expressing pilus adhesin RrgA, *Journal of Clinical Investigation* (2016). DOI: 10.1172/JCI84705

Provided by Karolinska Institutet

Citation: Special properties of pneumococci affect their ability to cause meningitis (2016, June 28) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2016-06-special-properties-pneumococci-affect-ability.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.