

Telehealth interventions provide limited health benefits, concludes study

June 2 2016



There is considerable optimism amongst policy-makers that greater use of digital health technologies ('telehealth') in combination with new ways of working could transform health care delivery, helping the NHS to be sustainable in the face of rising demand.

But in a study published today in the *British Medical Journal (BMJ)*,

researchers conclude that while telehealth could provide some benefit to some patients, it is not a simple answer to a complex problem.

Led by researchers from the University of Bristol's Centre for Academic Primary Care, and funded by the NIHR, the [Healthlines study](#) aimed to evaluate the effectiveness and cost-effectiveness of a new telehealth [service](#). The trial of the Healthlines Service involved participants receiving regular telephone calls from a health advisor, who supported them in addressing their own health goals and directed them to relevant online resources, or sent them the information by post.

The researchers recruited 641 adults with a high risk of heart disease or strokes (cardiovascular disease) to a randomised controlled trial of the Healthlines Service. Half the patients were given usual care while the other half were given usual care plus contact about once per month from a Healthlines Service advisor. They looked for improvements in the participants' risk of [cardiovascular disease](#), blood pressure, weight, body mass index, cholesterol and smoking status.

Professor Chris Salisbury, a GP and head of the Centre for Academic Primary Care, led the research. He said: "We found some evidence that the Healthlines Service led to modest improvement in overall cardiovascular risk for a minority of participants, but had no impact on average risk. But the Service did have other benefits, such as improvements in diet and physical activity, and patients were more satisfied with their access to care and the treatment they received."

In a separate paper, published recently in *The Lancet Psychiatry*, the researchers describe using the Healthlines Service to support patients with depression. Compared with usual care alone, intervention participants reported small improvements in depression and anxiety, better access to support and advice, greater satisfaction with the support they received, and improvements in self-management.

"There has been much enthusiasm about the potential of telehealth, but there have been very few studies testing a telehealth system in a real-world application on a large scale," said Professor Salisbury.

"The results of our studies show that while a telehealth intervention such as this can be beneficial to some patients, it is not the magic bullet that some policy-makers seem to think it is. However, it is a relatively low-cost intervention that can make health care more accessible for conditions that affect a large number of [patients](#), so there is good reason to continue exploring ways to make telehealth more effective in future."

The research team involved colleagues from the University of Sheffield, Bristol Randomised Trials Collaboration, NHS England South (Wessex), University of Southampton, Royal College of Surgeons in Ireland, Nottingham Clinical Trials Unit, Solent NHS Trust and NHS Direct.

The team is now analysing the cost-effectiveness of this intervention in detail, and hopes to publish its results before the end of the year.

More information: Chris Salisbury et al. Telehealth for patients at high risk of cardiovascular disease: pragmatic randomised controlled trial, *BMJ* (2016). [DOI: 10.1136/bmj.i2647](https://doi.org/10.1136/bmj.i2647)

Chris Salisbury et al. Effectiveness of an integrated telehealth service for patients with depression: a pragmatic randomised controlled trial of a complex intervention, *The Lancet Psychiatry* (2016). [DOI: 10.1016/S2215-0366\(16\)00083-3](https://doi.org/10.1016/S2215-0366(16)00083-3)

Chris Salisbury et al. Telehealth for patients at high risk of cardiovascular disease: pragmatic randomised controlled trial, *BMJ* (2016). [DOI: 10.1136/bmj.i2647](https://doi.org/10.1136/bmj.i2647)

Provided by University of Bristol

Citation: Telehealth interventions provide limited health benefits, concludes study (2016, June 2) retrieved 28 June 2024 from <https://medicalxpress.com/news/2016-06-telehealth-interventions-limited-health-benefits.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.