

## Brachial-ankle pulse wave velocity can predict mortality

July 28 2016



(HealthDay)—Brachial-ankle pulse wave velocity (baPWV) can predict



all-cause mortality in patients with diabetes after first non-traumatic lower extremity amputation (LEA), according to a study published online July 16 in the *Journal of Diabetes Investigation*.

Kazuki Ikura, M.D., from the Tokyo Women's Medical University School of Medicine, and colleagues conducted an observational historical cohort study of 102 Japanese patients with diabetes after a first non-traumatic LEA.

The researchers found that during a mean follow-up period of 3.3 years, 44 patients reached the end point of all-cause <u>mortality</u>. baPWV was a significant predictor for all-cause mortality in univariate and multivariate analyses (hazard ratios, 1.05 and 1.04, respectively; both P

"In conclusion, the present study may provide evidence that baPWV is a useful clinical predictor for all-cause mortality in <u>patients</u> with <u>diabetes</u> after first non-traumatic LEA," the authors write. "This finding should be confirmed in prospective studies with a larger sample size and a multi-center design."

More information: <u>Abstract</u> <u>Full Text</u>

Copyright © 2016 HealthDay. All rights reserved.

Citation: Brachial-ankle pulse wave velocity can predict mortality (2016, July 28) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2016-07-brachial-ankle-pulse-velocity-mortality.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.