

Study shows changes in brain activity after mindfulness therapy in adolescents

July 20 2016, by Alison Sampson

Anxiety disorders are among the most common psychiatric conditions affecting children and adolescents. While antidepressants are frequently used to treat youth with anxiety disorders, sometimes, antidepressants may be poorly tolerated in children who are at high risk of developing bipolar disorder.

Researchers at the University of Cincinnati (UC) are studying how [cognitive therapy](#) that uses mindfulness techniques, such as meditation, quiet reflection and facilitator-led discussion, may serve as an adjunct to pharmacological treatments.

The study published in the *Journal of Child and Adolescent Psychopharmacology*, looked at brain imaging in youth before and after mindfulness based therapy and saw changes in brain regions that control emotional processing. It is part of a larger study by co-principal investigators Melissa DelBello, MD, Dr. Stanley and Mickey Kaplan Professor and Chair of the UC Department of Psychiatry and Behavioral Neuroscience, and Sian Cotton, PhD, associate professor of family and community medicine, director of the UC's Center for Integrative Health and Wellness, looking at the effectiveness of mindfulness-based therapy.

In a small group of youth identified with [anxiety disorders](#) (generalized, social and/or separation anxiety) and who have a parent with bipolar disorder, researchers evaluated the neurophysiology of [mindfulness-based cognitive therapy](#) in children who are considered at-risk for developing bipolar disorder.

"Our preliminary observation that the mindfulness therapy increases activity in the part of the brain known as the cingulate, which processes cognitive and emotional information, is noteworthy," says Jeffrey Strawn, MD, associate professor in the Department of Psychiatry and Behavioral Neuroscience, director of the Anxiety Disorders Research Program and co-principal investigator on the study.

"This study, taken together with previous research, raises the possibility that treatment-related increases in brain activity [of the anterior cingulate cortex] during emotional processing may improve emotional processing in anxious youth who are at risk for developing [bipolar disorder](#)."

The study's findings in regard to increases in activity in the part of the brain known as the insula, the part of the brain responsible for monitoring and responding to the physiological condition of the body, are of high interest, Strawn adds.

In this pilot trial, nine participants ages 9 to 16 years, underwent functional magnetic resonance imaging (fMRI) while performing continuous performance tasks with emotional and neutral distractors prior to and following 12 weeks of mindful-based cognitive therapy.

"Mindfulness-based therapeutic interventions promote the use of meditative practices to increase present-moment awareness of conscious thoughts, feelings and body sensations in an effort to manage negative experiences more effectively," says Sian Cotton, PhD, an associate professor of family and community medicine at UC, director of the UC's Center for Integrative Health and Wellness and a co-author on the study. "These integrative approaches expand traditional treatments and offer new strategies for coping with psychological distress."

"Clinician-rated anxiety and youth-rated trait anxiety were significantly

reduced following treatment; the increases in mindfulness were associated with decreases in anxiety. Increasingly, patients and families are asking for additional therapeutic options, in addition to traditional medication-based treatments, that have proven effectiveness for improved symptom reduction. Mindfulness-based therapies for mood disorders is one such example with promising evidence being studied and implemented at UC."

"The path from an initial understanding of the effects of psychotherapy on brain activity to the identification of markers of treatment response is a challenging one, and will require additional studies of specific aspects of emotional processing circuits," says Strawn.

More information: Jeffrey R. Strawn et al, Neural Function Before and After Mindfulness-Based Cognitive Therapy in Anxious Adolescents at Risk for Developing Bipolar Disorder, *Journal of Child and Adolescent Psychopharmacology* (2016). [DOI: 10.1089/cap.2015.0054](https://doi.org/10.1089/cap.2015.0054)

Provided by University of Cincinnati Academic Health Center

Citation: Study shows changes in brain activity after mindfulness therapy in adolescents (2016, July 20) retrieved 20 March 2024 from <https://medicalxpress.com/news/2016-07-brain-mindfulness-therapy-adolescents.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--