

## **Counteracting poor decision-making due to sleep loss**

July 19 2016



The Washington State Sleep and Performance Research Center will develop and test cognitive flexibility training to combat the effects of sleep loss on decisionmaking under rapidly changing circumstances. Credit: Bob Hubner, Washington State University

Researchers from Washington State University's Sleep and Performance Research Center received a \$1.7 million grant to develop and test cognitive flexibility training to combat the effects of sleep loss on decision-making under rapidly changing circumstances.

Funding for the three-year project comes from the Congressionally Directed Medical Research Programs, a partnership between the U.S.



Congress, military and public to fund groundbreaking, high-impact medical research.

The project aims to reduce decision-making errors that contribute to failed military missions, industrial accidents, workplace injuries, financial losses and other serious consequences, said Hans Van Dongen, a research professor in WSU's Elson S. Floyd College of Medicine who is lead investigator.

"Sleep loss makes people very susceptible to poor decision-making, especially in dynamically changing environments such as military operations, emergency response scenarios and volatile financial markets," he said.

## **Building on earlier WSU findings**

Until recently, he said, the underlying brain mechanisms had been poorly understood and no meaningful solutions had been developed to address the problem.

The new project builds on studies by the WSU team that looked at the effects of <u>sleep</u> deprivation on the different cognitive processes involved in decision-making. Results suggested that sleep loss affects cognitive processes in different ways. For example, working memory was relatively unaffected by sleep deprivation, whereas sustained attention (the ability to focus on a task for an extended period of time) declined substantially.

One important discovery was that sleep deprivation causes people to have trouble acting on new information that signals changes in the situation they are dealing with. Although their brain perceives the information, it doesn't come through as important to the task at hand. This leads people to try to remedy the situation without changing their



behavior, even though their actions no longer have the desired effect.

The discovery of this "feedback blunting" phenomenon led the team to the idea of using cognitive flexibility training to help overcome the problem.

## **Cognitive flexibility despite sleep loss**

"If we can train people to have the cognitive flexibility to adapt to changing circumstances despite <u>sleep loss</u>, it would provide benefits to anyone holding a job that involves fast-moving circumstances and high stakes - from soldiers to surgeons and pilots to politicians," Van Dongen said. He added that cognitive flexibility training has been proven to be helpful under other circumstances but hasn't yet been tried in the context of sleep deprivation.

The WSU team - which includes psychology researchers Paul Whitney and John Hinson, from the College of Arts and Sciences, and Kimberly Honn and Matt Layton, College of Medicine faculty - will design a cognitive flexibility training program geared toward optimizing participants' situational awareness and minimizing decision errors.

## 90 will participate in study

The researchers will test the effectiveness of their training approach in a laboratory-based <u>sleep deprivation</u> experiment during which 60 study participants will be kept awake for 38 hours. Half will receive the cognitive flexibility training and half will not. Another 30 participants will form a control group that will receive the <u>cognitive flexibility</u> training but will not be sleep deprived.

Throughout the experiment, participants will complete computer-based



performance tasks specifically designed to measure cognitive functioning in high-pace, complex situations. Data from the experiment will be analyzed to determine whether those who were sleep deprived after receiving the training performed better than those who were sleep deprived but did not receive the training.

In addition, the research team will look at the extent to which the training is as effective in sleep-deprived participants as compared to the well-rested participants in the control group.

"With the solution we're proposing, we now - for the first time - have a fighting chance of getting a handle on this problem that the military and many others have been struggling with for so long," said Van Dongen.

Provided by Washington State University

Citation: Counteracting poor decision-making due to sleep loss (2016, July 19) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2016-07-counteracting-poor-decision-making-due-loss.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.