

3-D face scans may pinpoint autism earlier than normal

July 25 2016, by Tony Malkovic, Sciencenetwork Wa



Credit: AI-generated image ([disclaimer](#))

Telethon Kids Institute researchers are hoping 3-D face scanning technology will help detect autism in children as young as one year old, which is at least one year earlier than current practices allow for.

The study is scanning the [faces](#) of [children](#) with [autism spectrum](#)

[disorder](#) (ASD) as well as their siblings and parents to provide insights into ASD and earlier detection.

While autism isn't normally thought to have distinctive face features, recent studies found small facial anomalies.

"So we're recruiting children between the ages of nine and 15 months who have already been identified as showing warning signs and we're obtaining facial images from these children," Dr Gail Alvares, a researcher with the autism team at Telethon Kids Institute, says.

"We'll be following them up over the next few years to see which of those children will go on to develop autism."

One thing being examined is the so-called 'masculine' face.

"What we found is that boys who haven't been diagnosed with autism prior to puberty actually have more masculine faces," Dr Alvares says.

That is linked to testosterone exposure in the womb, according to the researchers.

"So the more testosterone you are exposed to as a foetus relates to how masculine your face is in your adult life, that and that's across males and females," she says.



3D face scans...autism researcher Dr Gail Alvares demonstrates the facial scanning equipment. Credit: Telethon Kids Institute

So is a masculine face an indicator of autism?

"Not necessarily, but there is this theory that higher levels of testosterone may have some effect on changing brain development in utero," she says.

Dr Alvares says [autism](#) can currently be reliably diagnosed in children as young as two by observing their behaviours.

"The problem is that we don't," she says.

"In Australia, we diagnose on average at about age four, which is far too late for early intervention to really have its best effect.

"Ideally, we would like to diagnose between 12 and 24 months.

"We know that children who have access to early and very intensive intervention before the age of three have better cognitive and language outcomes."

The Telethon researchers hope to compile a wider database of faces, including people who do not have ASD.

"So we'll be rolling this out at a bunch of community events and open days, so we'll be scanning lots of families as part of the Telethon weekend that's happening in October at the Perth Convention Centre," Dr Alvares says.

This article first appeared on [ScienceNetwork Western Australia](#) a science news website based at Scitech.

Provided by Science Network WA

Citation: 3-D face scans may pinpoint autism earlier than normal (2016, July 25) retrieved 19 April 2024 from <https://medicalxpress.com/news/2016-07-d-scans-autism-earlier.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--