


Diabetes impacts QOL outcome after lumbar decompression

July 6 2016



(HealthDay)—Diabetes predicts diminished quality of life (QOL)

improvements after lumbar decompression surgery, according to a study published in the June issue of *The Spine Journal*.

Michael P. Silverstein, M.D., from the Cleveland Clinic Center for Spine Health, and colleagues examined the effect of preoperative diabetes on postoperative change in QOL in a [retrospective cohort study](#) involving 212 [patients](#). Participants underwent lumbar decompression and were followed for a minimum of six months.

The researchers found that patients with diabetes experienced no significant changes in any QOL measures, while patients without diabetes experienced significant improvements in EuroQol 5-Dimensions (EQ-5D), Pain Disability Questionnaire, and Patient Health Questionnaire 9 (all P

"The burden of comorbidities may impact the QOL benefit of decompression spine surgery," the authors write. "In the present study, [diabetes](#) was found to independently predict diminished improvement in QOL after lumbar decompression."

Several authors disclosed financial ties to the medical technology industry.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Diabetes impacts QOL outcome after lumbar decompression (2016, July 6) retrieved 5 May 2024 from <https://medicalxpress.com/news/2016-07-diabetes-impacts-qol-outcome-lumbar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.