

# Gene plays role in poor speech processing, dyslexia

July 11 2016

---



The DCDC2 gene is required for normal auditory processing of complex speech, UT Dallas researchers found. Credit: Emw/Wikimedia Commons

A new study led by UT Dallas researchers shows that a gene associated with dyslexia may interfere with the processing of speech, ultimately leading to reading problems that are characteristic of the disorder.

While previous studies have tied the DCDC2 gene to [dyslexia](#), Dr. Michael Kilgard, a neuroscientist at UT Dallas and one of the study's co-

authors, said the new findings are the first to demonstrate that the gene is required for normal auditory processing of complex speech.

According to Kilgard, there are two theories about the cause of dyslexia: a visual or memory theory, and the auditory theory. "This study puts more weight on the side of the auditory theory," he said.

"We now have evidence that strongly suggests that people with dyslexia don't actually hear all of the sounds they need to hear," said Kilgard, who is the Margaret Fonde Jonsson Professor in the School of Behavioral and Brain Sciences and the associate director of the Texas Biomedical Device Center.

"If you have trouble hearing the sounds in your language, you will have trouble learning to read later," he said. "Armed with this information about a genetic link, we may be able to determine who is at risk for reading problems before they have trouble—before they even start learning to read."

The study, which appears in the *Journal of Neuroscience*, involved healthy rats that were taught a variety of speech sound discrimination tasks. When the DCDC2 gene was suppressed, the rats were virtually unable to identify specific speech sounds.

Kilgard equated the speech discrimination problems with issues experienced by children as they try to learn language skills.

He said the good news about the discovery is that individuals who have trouble distinguishing sounds could potentially recover that ability through therapy and neural training.

"Neural responses actually can be changed by training. Repetition does change those brain networks," Kilgard said. "The more you know about

genes, the more you can overcome the issues even if there is never a pill to solve the problem."

Kilgard is known for his research that uses vagus nerve stimulation paired with training to encourage neural plasticity, which is the reorganization of neural responses in the brain. He has shown that VNS can enhance plasticity and rehabilitation in multiple animal models and in patients with stroke and tinnitus.

Kilgard said the new findings provide a new tool that could change how we look at learning disabilities.

"Instead of making assumptions about what a child can achieve in life, children who have trouble distinguishing sounds can attain great things, although it may take more effort. Instead of getting there in 20 hours it may take 100 hours," he said.

Kilgard said certain commercial gene testing products can detect impaired genes, including DCDC2. But he said more work is needed to fully understand the link between neural processing and dyslexia.

**More information:** T. M. Centanni et al. Knockdown of Dyslexia-Gene Dcdc2 Interferes with Speech Sound Discrimination in Continuous Streams, *Journal of Neuroscience* (2016). [DOI: 10.1523/JNEUROSCI.4202-15.2016](https://doi.org/10.1523/JNEUROSCI.4202-15.2016)

Provided by University of Texas at Dallas

Citation: Gene plays role in poor speech processing, dyslexia (2016, July 11) retrieved 28 April 2024 from <https://medicalxpress.com/news/2016-07-gene-role-poor-speech-dyslexia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.