

Preventing HIV in transgender people—JAIDS assembles critical evidence

July 20 2016

Programs to reduce the high risk of HIV infection among transgender people are urgently needed—but efforts are hindered by a lack of accurate information on HIV prevalence, HIV incidence, and specific risk factors facing this key population. A special supplement to *JAIDS: Journal of Acquired Immune Deficiency Syndromes* presents essential information to meet the challenges of HIV prevention in the transgender population.

"The 'Transgender supplement' has been developed to fill critical gaps in information on the state of the HIV epidemic among [transgender](#) individuals, to discuss opportunities for culturally-tailored [prevention](#) interventions, and to inform the way forward in responding to the unique public health challenge in this often marginalized and underserved population," according to Guest Editors Kenneth Mayer of Fenway Health, Boston; Beatriz Grinsztejn of Fundação Oswaldo Cruz, Rio de Janeiro; and Wafaa M. El-Sadr of Columbia University, New York. The full contents of the special issue are freely available on the *JAIDS* website: <http://www.jaids.com>.

Experts Seek to Build Evidence for Effective HIV Prevention in Transgender People

Although transgender individuals account for less than one percent of the population, they have a "distinctively increased" burden of HIV disease. Recent estimates suggest that the worldwide prevalence of HIV

among transgender women is 19 percent—with odds HIV close to 50 times higher than in non-transgender adults of reproductive age.

Historically, HIV researchers and public health officials have grouped transgender women in the category of "men who have sex with men" (MSM). But that overlooks the multi-level factors—sexual behaviors, social networks, and discrimination, among others—that may contribute to HIV risk in transgender women. Even less is known about HIV risk and risk factors among [transgender men](#).

Knowledge gaps exist in other areas as well, including the effectiveness of pre-exposure prophylaxis (PrEP) using antiretroviral drugs to decrease HIV transmission. Social stigma and the threat of violence may prevent [transgender people](#) from accessing available testing, prevention, and treatment services. "Developing effective programs to reduce HIV in transgender individuals will require an in-depth understanding of the epidemiology of behaviors and risks with specific types of partners in the diverse cultures where transgender people live," according to Dr. Mayer and colleagues.

Building on a recent workshop sponsored by the [HIV Prevention Trials Network \(HPTN\)](#), the eight papers in the special issue frame the important issues in developing a more holistic approach to the engagement of HIV prevention interventions for transgender individuals. Topics include:

- A comprehensive analysis of data on the epidemiology of HIV in transgender people.
- Potential for interactions between antiretroviral drugs and hormone therapy used by some transgender individuals—a key factor in evaluating the use of PrEP.
- Initial data and research priorities on PrEP in transgender women.

- A "holistic framework" for understanding the health needs of transgender people.
- Behavioral approaches to improving HIV protection for transgender women.

More information: Kenneth H. Mayer et al. Transgender People and HIV Prevention, *JAIDS Journal of Acquired Immune Deficiency Syndromes* (2016). [DOI: 10.1097/QAI.0000000000001086](https://doi.org/10.1097/QAI.0000000000001086)

Provided by Wolters Kluwer Health

Citation: Preventing HIV in transgender people—JAIDS assembles critical evidence (2016, July 20) retrieved 3 May 2024 from <https://medicalxpress.com/news/2016-07-hiv-transgender-peoplejaids-critical-evidence.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--