

# Inhibiting ApoC-III reduces triglyceride levels in mice

July 11 2016

---

Some individuals remain at high risk for developing cardiovascular disease even after preventative treatment with cholesterol-lowering medications. This persistent risk has been linked to triglyceride levels, which can remain elevated in spite of substantially lowered LDL, the form of cholesterol associated with heart disease. Triglyceride concentrations are highly correlated to levels of apolipoprotein C-III (ApoC-III), suggesting that ApoC-III may be a potential target for controlling triglyceride levels.

In this month's issue of the *JCI*, a team led by Jeffrey Esko at the University of California San Diego determined that ApoC-III increases triglyceride levels in blood by interacting with the same pathways that control LDL cholesterol levels.

They found that treating mice with an inhibitor of ApoC-III reduced triglyceride levels, but this effect was prevented in mice lacking two key components of the LDL cholesterol signaling pathway.

These findings suggest that blocking ApoC-III may be an effective approach for reducing the risk of cardiovascular disease in patients with high [triglyceride levels](#).

**More information:** Philip L.S.M. Gordts et al, ApoC-III inhibits clearance of triglyceride-rich lipoproteins through LDL family receptors, *Journal of Clinical Investigation* (2016). [DOI: 10.1172/JCI86610](#)

Provided by JCI Journals

Citation: Inhibiting ApoC-III reduces triglyceride levels in mice (2016, July 11) retrieved 18 April 2024 from

<https://medicalxpress.com/news/2016-07-inhibiting-apoc-iii-triglyceride-mice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.