

Nonvascular thoracic MRI improves clinical decision making

July 28 2016



(HealthDay)—Assessment with nonvascular thoracic magnetic

resonance (MR) imaging improves clinical decision making, according to a study published in the August issue of *Radiology*.

Jeanne B. Ackman, M.D., from Massachusetts General Hospital in Boston, and colleagues examined the impact of nonvascular thoracic MR imaging on the clinical [decision making](#) and diagnostic certainty of thoracic surgeons. Seven thoracic surgeons participated in this two-year survey study after completing a one-time demographic survey. Each time a surgeon ordered a nonvascular thoracic MR imaging study they were sent a link to the pre-test survey, which assessed the clinical rationale for imaging and pre-test diagnostic certainty. The surgeon was then sent a link to a post-test [survey](#) upon completion of the MR imaging report. Data were included from 99 pre- and post-test surveys.

The researchers found that 65 percent of MR imaging studies were requested due to indeterminate findings on computed tomography. MR imaging correlated with a significant reduction in the number of planned surgical interventions (P

"In appropriate cases, assessment with nonvascular thoracic MR imaging substantially affects the [clinical decision](#) making and diagnostic certainty of [thoracic surgeons](#)," the authors write.

Several authors disclosed financial ties to the medical device industry.

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Nonvascular thoracic MRI improves clinical decision making (2016, July 28) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2016-07-nonvascular-thoracic-mri-clinical-decision.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.