

Pancreaticoduodenectomy costs high at safety-net hospitals

July 28 2016



(HealthDay)—Pancreaticoduodenectomy (PD) is associated with high

costs at safety-net hospitals, according to a study published online July 27 in *JAMA Surgery*.

Derek E. Go, from the University of Cincinnati School of Medicine, and colleagues modeled different techniques for reducing the cost of complex [surgery](#) performed at safety-net hospitals carrying out PD. Data were included for 15,090 [patients](#) who underwent PD during the five years of the study.

The researchers found that among safety-net hospitals, 28.0, 63.0, and 9.0 percent of patients were treated at low-burden hospitals (LBHs), medium-burden hospitals (MBHs), and high-burden hospitals (HBHs), respectively. Higher rates of complications or comorbidities were seen at HBHs, and they had more patients with increased illness severity. Perioperative mortality was 3.7 percent at HBHs, compared with 1.6 and 1.7 percent at LBHs and MBHs, respectively (P

"Redirecting patients away from safety-net hospitals for complex surgery may represent the best option for reducing [costs](#)," the authors write.

More information: [Abstract](#)

[Full Text \(subscription or payment may be required\)](#)

[Editorial \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Pancreaticoduodenectomy costs high at safety-net hospitals (2016, July 28) retrieved 10 April 2024 from

<https://medicalxpress.com/news/2016-07-pancreaticoduodenectomy-high-safety-net-hospitals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.