

Paths to autism: one or many?

July 19 2016

A new report in *Biological Psychiatry* reports that brain alterations in infants at risk for autism may be widespread and affect multiple systems, in contrast to the widely held assumption of impairment specifically in social brain networks.

Autism is diagnosed based on impairments in social and communication behaviors. These symptoms tend to emerge in the second year of life, but identifying abnormalities in early infancy could help researchers understand how autism develops and potentially allow clinicians to predict the disorder before it emerges.

Attempts to identify precursors have primarily focused on social behaviors, based on the assumption that abnormalities in social [brain](#) networks arise early in life and compound throughout development. But Dr. Mayada Elsabbagh from McGill University in Canada, and Dr. Mark Johnson, from Birkbeck, University of London, suggest that recent studies do not support the idea of a singular pathway in the development of autism.

In their synthesis of studies examining infants at risk for autism, Elsabbagh and Johnson highlight behavioral research providing evidence for general abnormalities during the first year of life. These include delayed motor maturation, higher level of perceptual sensitivity, and poor attention flexibility. The authors also highlight brain imaging studies that provide evidence for widespread alterations throughout [brain networks](#), rather than focal deficits in social networks.

The behavioral and imaging studies challenge the assumption of early social network abnormalities that persist throughout development and lead to emergence of the disorder.

"Our review reveals little support for localized deficits in social brain network systems within the first year of life," said Elsabbagh. "It instead favors the view that atypical development involving perceptual, attentional, motor, and social systems precede emerging autism and lead to overt behavioral symptoms by the second year."

The review suggests that focusing on a single deficit may not be sufficient to identify early warning signs and will likely adjust how researchers conceptualize the disorder.

"There has been a concerted effort to identify the final common neural pathways underlying symptoms and deficits for psychiatric disorders," said Dr. John Krystal, Editor of *Biological Psychiatry*. "Yet the perspective shared by Elsabbagh and Johnson suggests that there are widespread disturbances in brain development in [autism spectrum disorder](#) and that the prominent social deficits either reflect the fact that circuits underlying social behaviors are among the many circuits affected or that some functional deficits are emergent properties of multiple affected circuits."

More information: Mayada Elsabbagh et al. Autism and the Social Brain: The First-Year Puzzle, *Biological Psychiatry* (2016). [DOI: 10.1016/j.biopsych.2016.02.019](https://doi.org/10.1016/j.biopsych.2016.02.019)

Provided by Elsevier

Citation: Paths to autism: one or many? (2016, July 19) retrieved 5 May 2024 from

<https://medicalxpress.com/news/2016-07-paths-autism.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.