

Prevalence of diagnosed sleep disorders has risen among US veterans

July 15 2016

A new study found a six-fold increase in the age-adjusted prevalence of any sleep disorder diagnosis over an 11-year period among U.S. veterans. The largest increases were identified in patients with post-traumatic stress disorder (PTSD), other mental disorders, or combat experience. Results also show that the prevalence of PTSD tripled during the study period.

In a sample of more than 9.7 million U.S. veterans, the age-adjusted prevalence of [sleep disorders](#) increased from less than 1 percent in 2000 to nearly 6 percent in 2010. Sleep apnea was the most common [sleep](#) disorder diagnosis (47 percent) followed by insomnia (26 percent). Veterans with cardiovascular disease, cancer, or other chronic diseases also experienced higher rates of sleep disorder diagnoses relative to those without comorbid conditions.

Study results are published in the July issue of the journal *Sleep*.

"Veterans with PTSD had a very high sleep disorder prevalence of 16 percent, the highest among the various health conditions or other population characteristics that we examined," said Principal Investigator and senior author James Burch, PhD, Associate Professor in the Department of Epidemiology and Biostatistics in the Arnold School of Public Health at the University of South Carolina. "Because of the way this study was designed, this does not prove that PTSD caused the increase in sleep disorder diagnoses," noted Burch, who also is a Health Science Specialist at the WJB Dorn Department of Veterans Affairs

Medical Center in Columbia, South Carolina. "However, we recently completed a follow-up study, soon to be submitted for publication, that examined this issue in detail. In that study, a pre-existing history of PTSD was associated with an increased odds of sleep disorder onset."

According to the American Academy of Sleep Medicine, sleep apnea is a sleep-related breathing disorder characterized by abnormalities of respiration during sleep. The most common form of [sleep apnea](#) is [obstructive sleep apnea](#), which is characterized by repetitive episodes of complete or partial upper airway obstruction occurring during sleep. Insomnia involves a frequent and persistent difficulty initiating or maintaining sleep that results in general sleep dissatisfaction and daytime impairment.

The study population consisted of all U.S. veterans seeking care in the Veterans Health Administration system between FY2000 and FY2010. Of the total sample of 9,786,778 [veterans](#), 93 percent were men, and 751,502 were diagnosed with at least one sleep disorder.

Provided by American Academy of Sleep Medicine

Citation: Prevalence of diagnosed sleep disorders has risen among US veterans (2016, July 15) retrieved 28 April 2024 from

<https://medicalxpress.com/news/2016-07-prevalence-disorders-risen-veterans.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--