

Survival, surgical interventions for children with rare, genetic birth disorder

July 26 2016

Among children born with the chromosome disorders trisomy 13 or 18 in Ontario, Canada, early death was the most common outcome, but 10 percent to 13 percent survived for 10 years, according to a study appearing in the July 26 issue of *JAMA*. Among children who underwent surgical interventions, 1-year survival was high.

Trisomy 13 and 18 are genetic diagnoses associated with characteristic physical features and organ anomalies, often including cardiac malformations and neurologic impairments that occur in approximately 8 to 15 per 100,000 live births. Most [children](#) with these disorders die shortly after birth, although limited data suggest some children survive longer. Surgeries are controversial, and little evidence is available about outcomes. Lack of information about longer-term survival complicates clinical decision making.

Katherine E. Nelson, M.D., of the Hospital for Sick Children, Toronto, and colleagues examined survival and utilization of any type of surgery among children born in Ontario between April 1991 and March 2012 with a diagnosis code for trisomy 13 or 18 on a hospital record in the first year of life. All procedures classified as occurring in an operating room were categorized as major, intermediate, or minor surgeries.

The study included 174 children with trisomy 13 and 254 children with trisomy 18. Median survival times were 12.5 days for trisomy 13 and 9 days for trisomy 18. Average 1-year survival for trisomy 13 was 20 percent and 13 percent for trisomy 18. Ten-year survival for trisomy 13

was 13 percent and 10 percent for trisomy 18. Survival did not change over the study period. Forty-one children (24 percent) with trisomy 13 and 35 children (14 percent) with trisomy 18 underwent surgeries. Median age at first surgery for trisomy 13 was 92 days and for trisomy 18 it was 206 days. Analysis indicated 1-year survival after first surgery of 71 percent for trisomy 13 and 69 percent for trisomy 18.

The authors note that longer-term survival and use of [surgical interventions](#) were more common in this study than previously reported in population-based studies. They add that one factor likely contributing to this was the use of health administrative data from Ontario's single-payer health care system, which captures all surgical procedures and deaths.

More information: *JAMA*, [DOI: 10.1001/jama.2016.9819](https://doi.org/10.1001/jama.2016.9819)

Provided by The JAMA Network Journals

Citation: Survival, surgical interventions for children with rare, genetic birth disorder (2016, July 26) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-07-survival-surgical-interventions-children-rare.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--