

Study sheds light on true risk for pancreatic cancer in patients with cysts

July 11 2016

A look back at more than half a million patient records has established that patients with pancreatic cysts have a significantly higher overall risk of pancreatic cancer compared to those without such cysts, according to a study in the July issue of *GIE: Gastrointestinal Endoscopy*, the journal of the American Society for Gastrointestinal Endoscopy (ASGE).

Certain types of pancreatic cysts (such as mucinous neoplasms) are known to have malignant potential based on their form or structure. Surveillance (watching) or surgical resection (removal) are possible ways to manage these. But other types of cysts do not fit the "potentially malignant" profile at the time they are detected (described as "morphologically benign"). Because previous studies did not follow [patients](#) for more than two years, experts have not known what the overall long-term risk is for developing pancreatic cancer in patients with cysts.

The study, "Risk of pancreatic cancer in patients with [pancreatic cyst](#)," looked at the history of patients in the Veterans Health Administration database, examining all patient encounters from October 1998 through September 2007. They included 520,970 patients, 755 of whom were diagnosed with pancreatic cysts. The study did not differentiate among different types of pancreatic cysts.

After adjusting for age, race, smoking and alcohol use, which are independent risk factors for pancreatic cancer, the researchers followed the history of patients with and without pancreatic cysts from the time of

cyst diagnosis to the time of cancer diagnosis. Those who did not develop cancer were followed until the end of the study or until they were lost to follow-up.

In the group with cysts, 17 of the 775 patients were diagnosed with pancreatic cancer over the time period of the study. Of those, five were diagnosed with cancer within one to two years of the cyst diagnosis; six patients within two to four years; and three within four to six years. The remaining three patients were diagnosed with pancreatic cancer within six to eight years of their cyst diagnosis.

In the large group of patients without cysts, 1,206 out of 520,415 were eventually diagnosed with pancreatic cancer. The mean age of patients diagnosed across both groups was 62 years, 11.5 months.

The overall risk of pancreatic cancer was estimated to be 19.64 times greater in the group with pancreatic cysts.

Pancreatic cysts that have not produced symptoms are being detected more frequently in patients as a result of increased use and improved quality of abdominal imaging ordered for various reasons. Prevalence of pancreatic cysts as an incidental finding on [magnetic resonance imaging](#) (MRI) was reported to range from 2.4 percent to 19.6 percent in two studies from 2010. Surgery is performed in only a small fraction of patients with pancreatic cysts.

The authors noted the need for future studies to evaluate the risk of [pancreatic cancer](#) in patients with documented mucinous cysts, based on cyst size and patient age, to help create appropriate strategies for diagnostic evaluation and management.

More information: ASGE has published a related guideline in the July issue of *GIE: Gastrointestinal Endoscopy*, entitled "The role of endoscopy

in the diagnosis and treatment of cystic pancreatic neoplasms. The role of endoscopy in the diagnosis and treatment of cystic pancreatic neoplasms." [www.giejournal.org/article/S00 ... \(16\)30060-8/abstract](http://www.giejournal.org/article/S00... (16)30060-8/abstract)

Provided by American Society for Gastrointestinal Endoscopy

Citation: Study sheds light on true risk for pancreatic cancer in patients with cysts (2016, July 11)
retrieved 23 April 2024 from

<https://medicalxpress.com/news/2016-07-true-pancreatic-cancer-patients-cysts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.