

One week of bed rest lowers muscle mass, insulin sensitivity

July 13 2016



(HealthDay)—One week of bed rest is associated with a substantial

reduction in skeletal muscle mass and decreased whole-body insulin sensitivity, according to a study published online June 29 in *Diabetes*.

Marlou L. Dirks, from Maastricht University Medical Center in the Netherlands, and colleagues examined mechanisms underlying disuse-induced insulin resistance in a study involving 10 healthy young males who were subjected to one week of bed rest. Lean body mass and quadriceps cross-sectional area (CSA) were assessed before and after bed rest and VO_2 peak and leg strength were measured. Whole-body [insulin sensitivity](#) was also measured. In addition, [muscle biopsies](#) were obtained to assess muscle lipid fraction and markers of mitochondrial and vascular content.

The researchers observed a 1.4 ± 0.2 kg loss in lean tissue and a 3.2 ± 0.9 percent decrease in quadriceps CSA after bed rest (both P_2 peak (P

"One week of [bed rest](#) substantially reduces [skeletal muscle](#) mass and lowers whole-body insulin sensitivity, without affecting mechanisms implicated in high-fat diet-induced insulin resistance," the authors write.

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: One week of bed rest lowers muscle mass, insulin sensitivity (2016, July 13) retrieved 28 April 2024 from <https://medicalxpress.com/news/2016-07-week-bed-rest-lowers-muscle.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
