


CDC urges prevention, early recognition of sepsis

August 24 2016



(HealthDay)—Many cases of life-threatening sepsis could be recognized

and treated long before they cause severe illness or death, according to an Aug. 23 *Vital Signs* report from the U.S. Centers for Disease Control and Prevention.

Researchers from the CDC found that about 70 percent of patients with sepsis had used [health care services](#) recently or had chronic diseases that required regular [medical care](#).

Health care providers play a critical role in preventing sepsis, which includes following infection control measures such as hand washing and by ensuring patients get recommended vaccines, the CDC said. It's also essential to educate patients and their families about the need to prevent infections, manage [chronic health conditions](#), and seek immediate medical care if an infection doesn't improve.

The CDC had additional advice for [health care providers](#): Know the signs and symptoms of sepsis. If sepsis is suspected, order tests to determine if an infection is present, where it is, and what caused it. Start antibiotics and other recommended medical care immediately. Also, monitor patients closely and reassess antibiotic treatment within 24 to 48 hours or sooner to determine whether the type of antibiotics, dose, and duration are correct, or need to be changed.

More information: [More Information](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: CDC urges prevention, early recognition of sepsis (2016, August 24) retrieved 1 May 2024 from <https://medicalxpress.com/news/2016-08-cdc-urges-early-recognition-sepsis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.