

Researchers study how fluctuating hormones affect female vulnerability to cocaine addiction

August 3 2016



Linda Perrotti, UTA associate professor of psychology and principal investigator on this study. Credit: University of Texas at Arlington



Researchers at The University of Texas at Arlington are studying how fluctuating estrogen levels make females increasingly sensitive to the rewarding effects of cocaine and ultimately, vulnerable to cocaine addiction.

Repeated administrations of cocaine cause molecular neuroadaptations in the brain that trigger changes in behavior aimed towards obtaining more of the drug, primarily affecting the dopamine reward system. Longterm exposure to the drug can create permanent changes in brain circuitry and intracellular responses, causing the individual to maintain a neurobiological disposition to seeking drugs.

"Compared to men, <u>women</u> experience <u>higher levels</u> of craving and relapse during periods of abstinence and take larger amounts of cocaine during bouts of relapse," said Linda Perrotti, UTA associate professor of psychology and principal investigator on this study.

"Our study on hormonal effects could lead to customizable and differentiated addiction treatment and prevention measures for men, women, women on hormone-based birth control, post-menopausal women and women on <u>hormone replacement therapy</u>."

Perrotti received a three-year \$413,970 National Institutes of Health grant to fund bio molecular research on the effects of changing hormone levels on the principal molecules implicated in behavioral adaptations resulting from drug use.

The research team's preclinical work using rodent models demonstrated that females given hormone treatment preferred higher doses of cocaine. This finding suggests that estrogen may alter dopamine signaling and influence the strength of cocaine-associated cues.

"Estrogen also activates messenger pathways within the dopamine



reward system and we need to find out how it affects learning and memory circuits, making drugs more or less rewarding," Perrotti added. 'Does it signal to the brain, 'Try cocaine again, you liked it that first time?' And when there are higher levels of progesterone and less estrogen at another point in the hormonal cycle, is the female less likely to try again? All this has important implications for treatment."

Perry Fuchs, UTA chair of psychology, underlined the links of this research to the university's theme of health and the human condition, as one of the key pillars of UTA's <u>Strategic Plan 2020: Bold Solutions I</u> <u>Global Impact</u>.

"Historically, most biomedical models have been based on men but it is increasingly clear that what we know about addiction in men does not always hold up in women," Fuchs said. "Dr. Perrotti's research could provide essential insights to improve the quality of treatment given to women suffering from drug addiction."

The researchers will use rodent models to focus on three specific molecules associated with drug addiction and reward: extracellular signal-related kinase or ERK, which plays a critical role in cocaine reward and the consolidation of memories for cocaine associated cues; the FosB protein, which is thought to activate genes that increase sensitivity to the effects of the abused drug; and the response element binding protein CREB, which modulates cocaine reward and serves as a regulator of the negative emotional symptoms occurring during cocaine withdrawal.

Provided by University of Texas at Arlington

Citation: Researchers study how fluctuating hormones affect female vulnerability to cocaine addiction (2016, August 3) retrieved 7 May 2024 from https://medicalxpress.com/news/2016-08-fluctuating-hormones-affect-female-vulnerability.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.