

Gastric bypass surgery and alcohol sensitivity

August 8 2016, by From Mayo Clinic News Network, Mayo Clinic News Network

After undergoing gastric bypass surgery last year, having a glass of wine affects me much more than it used to. Is this typical?

A: Yes. In fact, the effects of [alcohol](#) are nearly doubled in people who have had [gastric bypass surgery](#), compared with those who haven't had the procedure.

Gastric bypass - one of the most common types of [bariatric surgery](#) in the U.S. - helps you reduce your [food intake](#) by creating a small gastric pouch. Before the surgery, food enters your stomach and passes into the [small intestine](#). After surgery, most of the stomach and the first part of your small intestine (duodenum) are bypassed, and a digestive route directs food into the middle section of your small intestine (jejunum). This helps you lose weight by limiting the amount of calories you can consume and absorb.

But, this direct route to your small intestine also allows your body to absorb alcohol more readily - and much faster. Your gastric pouch is unable to break down alcohol as effectively as your old stomach would have. In addition, your body weight is likely much lower than it was before your surgery, meaning you get a higher dose of alcohol per pound.

Several studies compared blood alcohol concentrations in women who'd undergone [gastric bypass](#) with those who hadn't had the surgery. The studies showed that, for women who'd had the operation, blood alcohol

concentrations peaked sooner and at approximately double the level of those who hadn't had the operation. Women who'd had the procedure also felt much more inebriated for longer.

To answer your question, increased sensitivity to alcohol is real. For you, having two drinks is the equivalent of having four drinks, at which point you're likely well above the legal limit. It's an important limitation to keep in mind for your safety and the safety of others. (adapted from Mayo Clinic Health Letter)

©2016 Mayo Foundation for Medical Education and Research
Distributed by Tribune Content Agency, LLC.

Citation: Gastric bypass surgery and alcohol sensitivity (2016, August 8) retrieved 27 April 2024 from <https://medicalxpress.com/news/2016-08-gastric-bypass-surgery-alcohol-sensitivity.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|