

HIV-infected adults with depression have increased risk for heart attack

August 24 2016

Among more than 26,000 human immunodeficiency virus (HIV)-infected adults, those with major depressive disorder (MDD) were more likely to experience a heart attack than those without MDD, according to a study published online by *JAMA Cardiology*.

With the advent of highly effective antiretroviral therapy and improved survival, people with HIV-infection are living longer and are now at an increased risk for cardiovascular disease (CVD). There is an urgent need to identify novel risk factors and primary prevention approaches for CVD in HIV. Although depression is prevalent in HIV-infected adults and is associated with future CVD in the general population, its association with CVD events has not been examined in the HIV-infected population.

Matthew S. Freiberg, M.D., M.Sc., of the Vanderbilt University School of Medicine, Nashville, Tenn., and colleagues conducted a study that included 26,144 HIV-infected veterans without CVD at baseline (1998-2003) participating in the U.S. Department of Veterans Affairs Veterans Aging Cohort Study from April 2003 through December 2009. At study entry, 4,853 [veterans](#) (19 percent) with [major depressive disorder](#) were identified.

The average age of those with MDD was 47 years and for those without MDD was 48 years. During 5.8 years of follow-up, 490 acute myocardial infarction (AMI; [heart attack](#)) events occurred. After adjustment for demographics, CVD risk factors, and HIV-specific factors, the

researchers found that HIV-infected adults with MDD had a 30 percent greater risk for having an AMI than did HIV-infected [adults](#) without MDD. This elevation in AMI risk was slightly lessened to 25 percent after further adjustment for other variables, such as hepatitis C infection, kidney disease, alcohol/cocaine abuse or dependence, and hemoglobin levels.

"Our findings raise the possibility that, similar to the general population, MDD may be independently associated with incident atherosclerotic CVD in the HIV-infected population. Considering the dearth of research in this area, future epidemiologic and mechanistic studies that include women and non-VA populations with HIV are needed," the authors write.

More information: *JAMA Cardiology*. Published online August 24, 2016; [DOI: 10.1001/jamacardio.2016.2716](https://doi.org/10.1001/jamacardio.2016.2716)

Provided by The JAMA Network Journals

Citation: HIV-infected adults with depression have increased risk for heart attack (2016, August 24) retrieved 6 May 2024 from <https://medicalxpress.com/news/2016-08-hiv-infected-adults-depression-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--