

# Radical treatment and examination combined can halve mortality from prostate cancer

August 8 2016

---

Men with very high-risk prostate cancer, who are treated at hospitals with a high proportion of administered radical local treatment (radiotherapy or prostatectomy), only have half of the mortality risk of men who are treated at hospitals with the lowest proportion. This is according to a new study conducted by researchers at Umeå University in Sweden and published in *European Urology*.

"The results suggest that the combination of thorough examination and subsequent radical local therapy can prolong life for men with very high-risk [prostate cancer](#)," says Pär Stattin, a researcher at the Department of Surgical and Perioperative Sciences who led the study.

Using patient data from the National Prostate Cancer Register (NPCR) of Sweden, the researchers investigated the link between prostatectomy or full-dose radiotherapy and mortality in men with deadly high-risk prostate cancer. Differences in treatment selection were analysed on a group basis whereas mortality from prostate cancer and other causes was analysed on an individual basis.

Results show that men with a very high-risk prostate cancer, who were treated at hospitals with the highest proportion of radical treatment, were only half as likely (mortality rate ratio; MRR 0.51) to die from prostate cancer compared to men treated at hospitals where radical local therapy was administered the least. However, since this was an observational study, the potential effects of high-intensity in diagnostic activity and work-up cannot be isolated from therapeutic activity.

"Our data suggests that there are benefits of radical local therapy for men with very advanced forms of prostate cancer. However, these [men](#) most often only receive hormonal treatment. To confirm our findings and hopefully to improve [treatment](#) praxis, there is a need for a randomized study of the effects of local radical therapy on patients with very high-risk prostate cancer," says Pär Stattin.

**More information:** Pär Stattin et al, Association of Radical Local Treatment with Mortality in Men with Very High-risk Prostate Cancer: A Semiecologic, Nationwide, Population-based Study, *European Urology* (2016). [DOI: 10.1016/j.eururo.2016.07.023](https://doi.org/10.1016/j.eururo.2016.07.023)

Provided by Umea University

Citation: Radical treatment and examination combined can halve mortality from prostate cancer (2016, August 8) retrieved 8 May 2024 from <https://medicalxpress.com/news/2016-08-radical-treatment-combined-halve-mortality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---