

A novel tool to predict septic kidney damage

August 4 2016



E coli bacteria. Credit: Fresh Science

Melbourne researcher Yugeesh Lankadeva has discovered what drives kidney damage and how to asses it through a patient's urine.

The new technique could warn doctors of septic kidney damage before it becomes life threatening.

Sepsis is increasingly common and is the leading cause of <u>acute kidney</u> <u>failure</u>, killing one in three people affected—3000 deaths each year in Australia.

Up until now, the mechanisms of <u>sepsis</u> were poorly understood, and there was no way to detect <u>kidney damage</u> before it was too late.



But Yugeesh, working at the Florey Institute of Neuroscience and Mental Health, has shown that real-time monitoring of <u>oxygen levels</u> in the urine of patients with sepsis could predict kidney failure before it happens, and save lives.

The new, non-invasive technique would allow doctors to pick up patients at risk of developing kidney failure and will be trialled in August 2016 at Melbourne's Austin Hospital.

Sepsis is caused by our own immune system over-reacting to an underlying infection. Around half of those patients then suffer kidney failure, usually within a day of developing sepsis. One third of those patients do not leave the hospital alive, because they are not diagnosed early enough for doctors to intervene.





Credit: Fresh Science

Sepsis can affect anyone and can be acquired both within our community (70 per cent of cases) and from hospitals and <u>health care</u> <u>facilities</u> (30 per cent). It is more prevalent amongst people with weakened immune systems, such as newborn infants and the elderly. It is



also a risk for patients with pre-existing health complications—such as cancer, diabetes AIDS and kidney or liver disease—who don't have the ability to fight off these infections.

The incidence of sepsis is increasing due to our aging population and the increasing number of drug-resistant infections.

But despite that, we didn't know exactly how kidneys fail in sepsis. And doctors could not detect sepsis until irreversible damage to the organs had already occurred.

Yugeesh was able to show that while sepsis is developing, more blood overall is sent towards the kidney, but less of this blood makes it to the kidney's inner zone. As that part of the organ becomes deprived of oxygen, he proposes that cells stop functioning normally, causing the kidneys to fail.

Because urine is produced in the kidneys, Yugeesh suspected that oxygen levels in the kidneys would be the same as those in urine. He measured urine oxygen using a probe inserted via a catheter up into the bladder, and he found he was right: oxygen levels in <u>urine</u> correlated really well with oxygen levels in the kidney's inner zone during development of septic kidney failure.

The new, real-time diagnostic tool can detect patients that are at risk of developing kidney failure before it happens, so that early preventative strategies can be implemented.

Yugeesh and colleagues will start testing this new <u>diagnostic tool</u> on patients in the Intensive Care Unit at the Austin Hospital in August 2016. It's already been shown to work in <u>patients</u> undergoing cardiac surgery, who also have a high risk of developing <u>kidney failure</u>.



Provided by Fresh Science

Citation: A novel tool to predict septic kidney damage (2016, August 4) retrieved 28 April 2024 from <u>https://medicalxpress.com/news/2016-08-tool-septic-kidney.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.