

# How brain separates relevant and irrelevant information

September 20 2016

---



Credit: public domain

Imagine yourself sitting in a noisy café trying to read. To focus on the book at hand, you need to ignore the surrounding chatter and clattering of cups, with your brain filtering out the irrelevant stimuli coming through your ears and "gating" in the relevant ones in your vision—words on a page.

In a new paper in the journal *Nature Communications*, New York University researchers offer a new theory, based on a [computational](#)

[model](#), on how the brain separates relevant from irrelevant information in these and other circumstances.

"It is critical to our everyday life that our brain processes the most important information out of everything presented to us," explains Xiao-Jing Wang, Global Professor of Neural Science at NYU and NYU Shanghai and the paper's senior author. "Within an extremely complicated [neural circuit](#) in the brain, there must be a gating mechanism to route relevant information to the right place at the right time."

The analysis focuses on inhibitory neurons—the brain's traffic cops that help ensure proper neurological responses to incoming stimuli by suppressing other neurons and working to balance excitatory neurons, which aim to stimulate neuronal activity.

"Our model uses a fundamental element of the brain circuit, involving multiple types of inhibitory neurons, to achieve this goal," Wang adds. "Our computational model shows that inhibitory neurons can enable a neural circuit to gate in specific pathways of information while filtering out the rest."

In their analysis, led by Guangyu Robert Yang, a doctoral candidate in Wang's lab, the researchers devised a model that maps out a more complicated role for inhibitory neurons than had previously been suggested.

Of particular interest to the team was a specific subtype of inhibitory neurons that targets the excitatory neurons' dendrites—components of a neuron where inputs from other neurons are located. These dendrite-targeting inhibitory neurons are labeled by a biological marker called somatostatin and can be studied selectively by experimentalists. The researchers proposed that they not only control the overall inputs to a

neuron, but also the inputs from individual pathways—for example, the visual or auditory pathways converging onto a neuron.

"This was thought to be difficult because the connections from inhibitory neurons to excitatory neurons appeared dense and unstructured," observes Yang. "Thus a surprising finding from our study is that the precision required for pathway-specific gating can be realized by [inhibitory neurons](#)."

The study's authors used computational models to show that even with the seemingly random connections, these dendrite-targeting neurons can gate individual pathways by aligning with excitatory inputs through different pathways. They showed that this alignment can be realized through synaptic plasticity—a brain mechanism for learning through experience.

Provided by New York University

Citation: How brain separates relevant and irrelevant information (2016, September 20) retrieved 1 May 2024 from <https://medicalxpress.com/news/2016-09-brain-relevant-irrelevant.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--