

# Low cancer symptom awareness linked to lower chance of survival

September 28 2016

---

In regions where cancer survival is poorer, people on average have lower awareness of cancer symptoms, according to new findings published today (Wednesday) in the *British Journal of Cancer*.

Researchers also found that general awareness of cancer symptoms varies greatly across England.

This is the first study in England to look at whether regional variation in cancer survival might be linked to [regional variation](#) in cancer symptom awareness and barriers to seeing the GP.

On average, each additional cancer symptom recognised was associated with about a 1.6 per cent increase in 1 year overall cancer survival for patients in that region.

In this study, only breast cancer survival was associated with perceived barriers to seeing the doctor - such as embarrassment and difficulties getting transport to the doctor's surgery.

East London had the lowest awareness of cancer symptoms and highest average number of barriers to seeking medical help among the region's residents. The highest cancer awareness was found in areas such as Peterborough, Bedfordshire and Cambridgeshire.

Dr Maja Niksic, lead author at the London School of Hygiene & Hygiene Tropical Medicine, said: "A range of things can influence

cancer survival. Stage of disease at diagnosis is one of them, and if we can ensure more people know what to look out for and see their GP if they notice any unusual or persistent changes, we may be able to reduce the numbers of patients who are diagnosed with advanced disease, where curative treatment is not often an option.

"Based on our research we think that health campaigns should focus on helping people to recognise cancer symptoms early and seek medical advice about these – especially in socio-economically deprived areas, where cancer survival is generally lower."

"Research like this helps reinforce the value of the Government's investment in public cancer awareness campaigns, and can be used to better inform and direct efforts."

Jessica Kirby, senior health information manager at Cancer Research UK, said: "It's important the public are informed about getting to know their body and what's normal for them, and spotting potential cancer signs and symptoms. And we need to ensure people feel they can go and see their GP if they notice any unusual changes.

"Many things influence [cancer survival](#) and there could be other explanations for the variation in survival across the UK. For example, risk factors like smoking could influence survival, and the way health services are set up may also play a part. The NHS should ensure that everyone can have the right information and access to services to give them the best chance of being diagnosed and treated promptly, wherever they live."

**More information:** Maja Niksic et al. Is cancer survival associated with cancer symptom awareness and barriers to seeking medical help in England? An ecological study, *British Journal of Cancer* (2016). [DOI: 10.1038/bjc.2016.246](https://doi.org/10.1038/bjc.2016.246)

Provided by Cancer Research UK

Citation: Low cancer symptom awareness linked to lower chance of survival (2016, September 28) retrieved 1 May 2024 from <https://medicalxpress.com/news/2016-09-cancer-symptom-awareness-linked-chance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.