

Despite chronic pain, South Africans with HIV-related pain are surprisingly active

September 13 2016



Credit: Wits University

When one thinks about chronic conditions that are commonly painful, HIV doesn't typically spring to mind. However, more than 50% of HIV-positive individuals experience a painful condition like headache, chest

pain or neuropathy, and that pain is frequently experienced as moderate to severe in intensity.

What struck researchers from the Brain Function Research Group (BFRG) at Wits University as odd was that despite this high burden of pain in HIV, a couple of papers have emerged suggesting that, having asked patients, functional interference (having difficulty with things like walking or going to work) was not as great as they might have expected. One of these papers was from the BFRG and had been completed locally in Johannesburg, South Africa.v

To investigate whether pain does actually affect function in HIV (as it does in many other clinical conditions), researchers Dr Antonia Wadley, Emeritus Professor Duncan Mitchell and Associate Professor Peter Kamerman from the BFRG, based in the School of Physiology, Faculty of Health Sciences at Wits, conducted a cross-sectional study.

The results from the study, titled: Resilience does not explain the dissociation between chronic pain and physical activity in South Africans living with HIV, are published today, 13 September 2016, in the journal *PeerJ*.

To explain why pain may not affect function, the researchers first put it down to African patients being resilient, i.e. the ability to cope with adversity.

Explains Wadley: "No one has assessed resilience in people living with HIV and chronic pain before. We hypothesised that people living with HIV would be generally be pretty resilient, and those who were more resilient would be more active and report lower pain intensity."

Measuring resilience in HIV-patients objectively for

the first time

For the study, the researchers recruited HIV-positive patients from an HIV clinic in Johannesburg: half with chronic pain (defined as having had pain most days for at least three months) and half without. They then assessed resilience and, as well as asking patients about their activity, the researchers measured it objectively for the first time in a subset of patients using accelerometers, which are like sophisticated pedometers. They also asked the patients about their day-to-day worries.

"It turns out," Wadley says, "that we were right on one thing, HIV-positive patients in our study were really resilient, but our hypothesis was wrong: being more resilient did not associate with being more active or having lower pain intensity. In fact, the activity results astounded us. Not only was patients' activity not as affected as one might expect, it wasn't affected at all."

There was no difference in activity intensity, duration, or time spent at different intensities of activity between those with and without chronic pain. "This is something you just don't see in other types of long term pain," she adds.

The researchers then looked at how frequently patients worried about their health, money, food and family. "We thought that if patients were worried about money and having enough food, that pain might be relegated to a lower priority." They found that patients in [chronic pain](#) worried more frequently about each of these things compared to their pain-free counterparts and that health was lowest down the list.

"So it really does appear that if you are poor, pain may be relegated to a lower priority. Indeed, our analysis showed that worrying more about food associated with higher levels of activity," says Wadley.

Going forward

The researchers also asked the patients in pain what else they worried about and who they had told about their pain. "It turns out that HIV-related stigma is a real problem and that half the [patients](#) in pain had not told their closest friends and some not even their family about their pain, for fear that it might reveal their HIV status."

Wadley says it thus seems that economic stresses and fear of HIV-related stigma may drive people to maintain high levels of activity, even when they are in severe pain.

"What's not clear is whether this kind level of activity in the face of [pain](#) is helpful or harmful and that's something we will be looking into next," she adds.

Provided by Wits University

Citation: Despite chronic pain, South Africans with HIV-related pain are surprisingly active (2016, September 13) retrieved 20 March 2024 from <https://medicalxpress.com/news/2016-09-chronic-pain-south-africans-hiv-related.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
