

# Drop in use of digital rectal examination, PSA testing

September 20 2016

---



(HealthDay)—Following U.S. Preventive Services Task Force

(USPSTF) recommendations against routine prostate-specific antigen (PSA) screening, there has been a decrease in utilization of digital rectal examination and PSA testing, according to a study published in the October issue of *The Journal of Urology*.

Jonathan Shoag, M.D., from New York Presbyterian Hospital in New York City, and colleagues characterized trends in the rate of digital rectal examination and PSA testing by [primary care physicians](#) using the National Ambulatory Medical Care Survey for men older than 40 years presenting for preventive care. Data were included for 3,368 visits for the study of digital rectal examination trends from 2005 to 2012, and for 4,035 visits for the study of PSA trends from 2002 to 2012.

The researchers observed a decrease in the proportion of visits where digital rectal examination was performed following the USPSTF recommendation, from 16.0 to 5.8 percent (64 percent decrease; P

"This suggests that [prostate cancer screening](#) is rapidly disappearing from primary care practice," the authors write.

**More information:** [Abstract](#)  
[Full Text](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Drop in use of digital rectal examination, PSA testing (2016, September 20) retrieved 2 May 2024 from <https://medicalxpress.com/news/2016-09-digital-rectal-psa.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--