

Early TIPS beneficial in acute esophageal variceal bleeding

September 26 2016



(HealthDay)—For patients presenting with acute esophageal variceal

bleeding (EVB), early transjugular intrahepatic portosystemic shunt (TIPS) is associated with reductions in in-hospital rebleeding and mortality, with no increase in hepatic encephalopathy, according to a study published online Sept. 14 in the *Journal of Gastroenterology and Hepatology*.

Basile Njei, M.D., M.P.H., from the Yale University School of Medicine in New Haven, Conn., and colleagues examined the impact of early TIPS on outcomes for U.S. patients hospitalized with EVB from 2000 and 2010. Data were included for 142,539 patients with EVB and decompensated cirrhosis.

The researchers observed a decrease in the age-adjusted in-hospital mortality rate from 656 to 412 per 100,000 (37.2 percent decrease; P

"Early preventive TIPS in patients with EVB and decompensated cirrhosis was associated with significant in-hospital reductions in rebleeding and mortality without a significant increase in encephalopathy in 'real-world' U.S. clinical practice," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Early TIPS beneficial in acute esophageal variceal bleeding (2016, September 26)
retrieved 5 May 2024 from
<https://medicalxpress.com/news/2016-09-early-beneficial-acute-esophageal-variceal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
