

Study explores link between weight and stroke risk in women

September 8 2016



(HealthDay)—Excess weight may put women at increased risk for

ischemic stroke, but at lower risk for hemorrhagic stroke, according to research published online Sept. 7 in *Neurology*.

The study was led by Gillian Reeves, Ph.D., of the University of Oxford in the United Kingdom. Her team tracked the 12-year health history of British women who averaged 57 years of age.

During that time, rates of [ischemic stroke](#) were 1.0 percent among [obese women](#) and 0.7 percent among women with a healthy weight. Rates of [hemorrhagic stroke](#) were 0.4 percent among obese women and 0.5 percent among women with a healthy weight.

"In U.K. women, higher body mass index is associated with increased risk of ischemic stroke but decreased risk of hemorrhagic stroke," the authors write. "The totality of the available published evidence suggests that body mass index-associated risks are greater for ischemic than for hemorrhagic stroke."

More information: [Abstract](#)
[Full Text](#)
[Editorial \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Study explores link between weight and stroke risk in women (2016, September 8) retrieved 26 April 2024 from <https://medicalxpress.com/news/2016-09-explores-link-weight-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.