

# Infrared light to detect early signs of esophageal cancer

September 6 2016

---

Scientists have developed an endoscope that uses near-infrared light to spot early warning signs of oesophageal - food pipe - cancer, according to research published in the *Journal of Biomedical Optics* today.

Researchers at the Cancer Research UK Cambridge Institute sprayed a [dye](#) on oesophageal tissue samples taken from people with Barrett's oesophagus - a condition that increases the risk of developing [oesophageal cancer](#). The dye sticks to healthy oesophageal cells but not to pre-cancerous cells.

They then shone [near-infrared light](#) on the tissue, which is just beyond the red colours that our eyes can normally see, and used a special camera to detect the near-infrared light the dye gave out.

Researchers had tried this approach before but had used a dye that gave out visible light to spot pre-cancerous cells. Problems arose because when cells are exposed to visible light, they naturally emit [visible light](#) themselves - making the distinction between healthy and abnormal cells hard to see.

But, by using a dye that gives out near-infrared light, scientists refined the process so they could make the distinction.

This new technique could be used to monitor people with Barrett's oesophagus who have an increased risk of developing oesophageal [cancer](#), and spot early signs that cancer might be developing before the

person has any symptoms.

Dr Sarah Bohndiek, scientist at the Cancer Research UK Cambridge Institute, said: "This research gives us hope for developing better techniques to detect the disease early.

"People who are at a high risk of developing oesophageal cancer, such as those with Barrett's oesophagus, could be closely monitored with this technique. And removing patches of pre-cancerous [cells](#) could prevent some cases of oesophageal cancer. But we need to do some further testing before clinical trials with patients can be set up to see how effective the approach could be at saving lives."

There are around 7,800 deaths from oesophageal cancer every year in the UK. For men, oesophageal cancer is the fourth most common cause of cancer death.

Dr Emma Smith, science communication manager at Cancer Research UK, said: "Early detection is crucial if we want to cure more cancers and save more lives. And oesophageal cancer is often caught at an advanced stage, meaning survival rates of the disease have remained stubbornly low.

"We urgently need new ways to detect the disease early. This development could lead to a cheaper, better way of detecting the [warning signs](#) of oesophageal cancer and even stop some people developing the disease."

**More information:** Dale J. Waterhouse ; James Joseph ; André A. Neves ; Massimiliano di Pietro ; Kevin M. Brindle ; Rebecca C. Fitzgerald ; Sarah E. Bohndiek. Design and validation of a near-infrared fluorescence endoscope for detection of early esophageal malignancy. *Journal of Biomedical Optics*. [DOI: 10.1117/1.JBO.21.8.08400](https://doi.org/10.1117/1.JBO.21.8.08400)

Provided by Cancer Research UK

Citation: Infrared light to detect early signs of esophageal cancer (2016, September 6) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2016-09-infrared-early-esophageal-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.