

Kids with bipolar disorder more likely to abuse drugs, alcohol: study

September 16 2016



(HealthDay)—For some teens with bipolar disorder, the risk that they

will abuse alcohol and drugs may increase as they get older, a new study suggests.

The research included 105 young people with bipolar disorder and 98 without the illness (the "control" group). Their average age was 14 when they first enrolled in the study. Bipolar disorder causes unusual shifts in mood, energy and activity levels, and also affects the ability to carry out everyday tasks.

Initially, 34 percent of the young teens with bipolar disorder also had "[substance use disorder](#)," which means they had a problem with drug or alcohol abuse. Only 4 percent of the kids in the control group abused alcohol or drugs, the study showed.

In addition, the researchers found, almost one-quarter of kids with bipolar disorder smoked cigarettes, compared to just 4 percent of the comparison group.

Five years after the initial study, the investigators followed up with 68 of the original bipolar patients and 81 members of the [control group](#). Among these participants, almost half of those with bipolar disorder had substance use disorder. Only 26 percent of those without the [mental health disorder](#) had trouble with drugs or drinking, the study revealed.

The risk was particularly high for people with persistent bipolar symptoms, the study authors said.

Dr. Timothy Wilens was the study's lead researcher. He's chief of child and adolescent psychiatry at MassGeneral Hospital for Children in Boston. He explained that bipolar disorder symptoms usually show up before kids start using drugs, cigarettes or alcohol.

That means health-care professionals who treat children with bipolar

disorder should also look for signs of substance abuse or cigarette smoking, Wilens said in a hospital news release.

The study also looked at the combination of bipolar disorder and [conduct disorder](#). Conduct disorder is a condition characterized by defiant, impulsive and possibly criminal behavior, according to the U.S. National Library of Medicine. When the two disorders were combined, the risk of smoking and alcohol and drug abuse was even greater, the study found.

Attention-deficit/hyperactivity disorder (ADHD), however, didn't seem to have an impact on the risk of drug and [alcohol abuse](#) in people with bipolar disorder.

"We were surprised to find that conduct disorder, but not ADHD, played such a large role in [increasing the risk] of substance use disorder," Wilens said.

He said it's possible that conduct disorder is at the root of the drinking and drug use as teens with [bipolar disorder](#) become young adults.

Findings from the study were published recently in the *Journal of Clinical Psychiatry*.

More information: The U.S. National Institute of Mental Health has more on [bipolar disorder](#).

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Kids with bipolar disorder more likely to abuse drugs, alcohol: study (2016, September 16) retrieved 23 April 2024 from <https://medicalxpress.com/news/2016-09-kids-bipolar-disorder-abuse-drugs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.