


Researchers study mild electrical stimulation for schizophrenia patients

September 29 2016



Functional magnetic resonance imaging (fMRI) and other brain imaging technologies allow for the study of differences in brain activity in people diagnosed with schizophrenia. The image shows two levels of the brain, with areas that were more active in healthy controls than in schizophrenia patients shown in orange, during an fMRI study of working memory. Credit: Kim J, Matthews NL, Park S./PLoS One.

A safe, noninvasive weak electrical current delivery called transcranial direct current stimulation (tDCS) will be investigated for its potential in treating schizophrenia patients in a study led by Raymond Cho, M.D., M.Sc., at McGovern Medical School at The University of Texas Health

Science Center at Houston (UTHealth).

The study, which comes with a two-year, \$850,000 grant from the National Institute of Mental Health (R61MH110044), will seek to answer whether tDCS can affect cognitive impairments in people with schizophrenia.

"Cognitive deficits such as [cognitive control](#) and working memory are among the most debilitating aspects of the illness and poorly treated by current medications," said Cho, associate professor in the Department of Psychiatry and Behavioral Sciences and director of the Clinical Cognitive Neuroscience Laboratory at McGovern Medical School at UTHealth. "If you have cognitive impairments, daily living – doing chores, going to school, working, driving – can be very challenging."

In the treatment, a weak electrical current is passed through the [dorsolateral prefrontal cortex](#) area of the brain between two electrodes placed on the scalp. In a previous study published in 2012 in the *American Journal of Psychiatry*, using tDCS on that same area of the brain was shown to reduce auditory hallucinations in [schizophrenia patients](#).

"The theory is that the electric current will increase excitability of the brain tissue, which will help that part of the brain function better," Cho said.

Patients remain fully conscious during the treatments, which do not require anesthesia or other medications, and are done in 20-minute periods twice a day for five days. Cho said patients generally feel nothing or a mild tingling when the therapy begins.

Researchers seek to determine the best dose of tDCS in the randomized study. Up to 100 patients will be enrolled. They will receive either a dose

of the therapy or a placebo. Patients will undergo cognitive testing and imaging before and after treatment. Find

Provided by University of Texas Health Science Center at Houston

Citation: Researchers study mild electrical stimulation for schizophrenia patients (2016, September 29) retrieved 18 April 2024 from <https://medicalxpress.com/news/2016-09-mild-electrical-schizophrenia-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.