

# Shorter duration of feeding tube placement with IMRT

September 30 2016

---



(HealthDay)—For patients with head and neck cancer (HNC), the

duration of feeding tube placement is shorter for those who receive definitive intensity-modulated radiation therapy (IMRT) compared with those who receive three-dimensional radiation therapy (3DRT), according to a study published online Sept. 23 in *Cancer*.

Beth M. Beadle, M.D., Ph.D., from the University of Texas MD Anderson Cancer Center in Houston, and colleagues used the linked Surveillance, Epidemiology, and End Results-Medicare database to determine the cohort, demographics, and cancer-related variables for patients with HNC. Treatment details were analyzed using claims data.

The researchers identified 2,993 patients. Overall, 54.4 percent of patients had ever had a [feeding tube](#) placed at a median follow-up of 47 months, with a median duration of 277 days from feeding tube placement to removal. The rates of feeding tube placement were similar for patients who received IMRT and 3DRT (odds ratio, 1.10; P = 0.35); patients who received 3DRT had a feeding tube in place for significantly longer than those who received IMRT (1.18 times longer; P = 0.03). The difference was only seen for patients who received definitive RT; no significant difference was seen in feeding tube placement or duration for patients who underwent surgery and received adjuvant RT.

"These data suggest that there may be significant quality-of-life benefits to IMRT with respect to long-term swallowing function in [patients](#) with HNC," the authors write.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Shorter duration of feeding tube placement with IMRT (2016, September 30) retrieved

23 April 2024 from

<https://medicalxpress.com/news/2016-09-shorter-duration-tube-placement-imrt.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.